

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 February 2002 (07.02.2002)

PCT

(10) International Publication Number  
**WO 02/10187 A1**

(51) International Patent Classification<sup>7</sup>: C07H 21/04,  
C12N 15/00, 15/12, 15/63, C07K 16/46

(74) Agent: MACPHAIL, Stuart; Fish & Richardson P.C., 45  
Rockefeller Plaza, Suite 2800, New York, NY 10111 (US).

(21) International Application Number: PCT/US01/41430

(22) International Filing Date: 26 July 2001 (26.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/220,991 27 July 2000 (27.07.2000) US

(63) Related by continuation (CON) or continuation-in-part  
(CIP) to earlier application:  
US 60/220,991 (CON)  
Filed on 27 July 2000 (27.07.2000)

(81) Designated States (*national*): AF, AG, AI, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GR, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,  
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).

(71) Applicant (*for all designated States except US*): MAYO  
FOUNDATION FOR MEDICAL EDUCATION AND  
RESEARCH [US/US]; 200 First Street S.W., Rochester,  
MN 55905 (US).

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): CHEN, Lieping  
[US/US]; 2821 Char Lane, N.E., Rochester, MN 55906  
(US).

Published:

— with international search report  
— before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

WO 02/10187 A1

(54) Title: B7-H3 AND B7-H4, NOVEL IMMUNOREGULATORY MOLECULES

(57) Abstract: The invention provides novel B7-H3 and B7-H4 polypeptides useful for co-stimulating T cells, isolated nucleic acid molecules encoding them, vectors containing the nucleic acid molecules, and cells containing the vectors. Also included are methods of making and using these co-stimulatory polypeptides.

## B7-H3 AND B7-H4, NOVEL IMMUNOREGULATORY MOLECULES

### BACKGROUND OF THE INVENTION

The invention is generally in the field of immunoregulation, and specifically T  
5 cell response regulation.

Mammalian T lymphocytes recognize antigenic peptides bound to major  
histocompatibility complex (MHC) molecules on the surface of antigen presenting  
cells (APC). The antigenic peptides are generated by proteolytic degradation of  
protein antigens within the APC. The interaction of the T cells with the APC and the  
10 subsequent response of the T cells are qualitatively and quantitatively regulated by  
interactions between cell surface receptors on the T cells with both soluble mediators  
and ligands on the surface of APC.

### SUMMARY OF THE INVENTION

15 The invention is based on the cloning of three human cDNA molecules  
encoding three novel polypeptides that co-stimulate T cell responses and on the  
functional characterization of the polypeptides that the cDNA molecules encode.  
Two of these novel co-stimulatory polypeptides, the B7-H3 polypeptides, differ from  
each other by a single amino acid residue (position number 166). The B7-H3  
20 polypeptide with serine at position 166 is designated herein B7-H3.1 (SEQ ID NO:1)  
and that with proline at position 166 is designated B7-H3.2 (SEQ ID NO:3). Text that  
refers to B7-H3 without specifying B7-H3.1 or B7-H3.2 is pertinent to both  
polypeptides. The third novel co-stimulatory polypeptide, B7-H4, is encoded by a  
separate gene. The invention features DNA molecules encoding the B7-H3 and B7-  
25 H4 polypeptides, functional fragments of the polypeptides, and fusion proteins  
containing the polypeptides or functional fragments of the polypeptides, B7-H3 and  
B7-H4 and functional fragments of both, vectors containing the DNA molecules, and  
cells containing the vectors. Also included in the invention are antibodies that bind to  
the B7-H3 and B7-H4 polypeptides. The invention features *in vitro*, *in vivo*, and *ex*  
30 *vivo* methods of co-stimulating T cell responses, methods of screening for compounds

that inhibit or enhance T cell responses, and methods for producing the polypeptides and fusion proteins.

Specifically the invention features an isolated DNA including: (a) a nucleic acid sequence that (i) encodes a polypeptide with the ability to co-stimulate a T cell, and (ii) hybridizes under stringent conditions to the complement of a sequence that encodes a polypeptide with an amino acid sequence with SEQ ID NO:1, or SEQ ID NO:3, or SEQ ID NO:5; or (b) a complement of this nucleic acid sequence. The polypeptide-encoding nucleic acid sequence included in the isolated DNA will be at least 10 bp, 15 bp, 25 bp, 50 bp, 75 bp, 100 bp, 125 bp, 150 bp, 175 bp, 200 bp, 250 bp, 300 bp, 350 bp, 400 bp, 450 bp, 500 bp, 550 bp, 600 bp, 650 bp, 700 bp, 750 bp, 800 bp, 840 bp, 850 bp, 900 bp, or 940 bp long. The nucleic acid sequence can encode a B7-H3 polypeptide that includes an amino acid sequence (a) extending from amino acid 27, 28, 29, 30, 31, 32, 33, or 34 to amino acid 316 of either SEQ ID NO:1 or SEQ ID NO:3, or (b) of SEQ ID NO:1 or SEQ ID NO:3. Alternatively, the nucleic acid sequence can have a nucleotide sequence with SEQ ID NO:2 or SEQ ID NO:4. The nucleic acid sequence can encode a B7-H4 polypeptide that includes an amino acid sequence with SEQ ID NO:5 or it can have a nucleotide sequence with SEQ ID NO:6. The nucleic acid sequence can also encode functional fragments of these B7-H3 or B7-H4 polypeptides.

The invention also embodies an isolated B7-H3 polypeptide encoded by a DNA that includes a nucleic acid sequence that (i) encodes a polypeptide with the ability to co-stimulate a T cell and (ii) hybridizes under stringent conditions to the complement of a sequence that encodes a polypeptide with an amino acid sequence with SEQ ID NO:1 or SEQ ID NO:3. The B7-H3 polypeptide can include an amino acid sequence starting at residue 27, 28, 29, 30, 31, 32, 33 or 34 and extending to 361 of SEQ ID NO:1 or SEQ ID NO:3. The invention also encompasses B7-H3 polypeptides that include an amino acid sequence with SEQ ID NO:1 or SEQ ID NO:3, or either of these amino acid sequences but differing solely by one or more (e.g., two, three, four, five, six, seven, eight, nine, ten, 12, 15, 20, 25, 30, 35, 40, 50, 60, 80, or 100) conservative substitutions.

The invention also includes an isolated B7-H4 polypeptide encoded by DNA that includes a nucleic acid sequence that (i) encodes a polypeptide with the ability to co-stimulate a T cell and (ii) hybridizes under stringent conditions to the complement of a sequence that encodes a polypeptide with an acid sequence with SEQ ID NOS:5.

- 5 The B7-H4 polypeptide can include an amino acid sequence starting at residue 27, 28, 29, 30, 31, 32, 33, or 34 and extending to residue 282 of SEQ ID NO: 5 or this amino acid sequence but differing solely by one or more (e.g., two, three, four, five, six, seven, eight, nine, ten, 12, 15, 20, 25, 30, 35, 40, 50, 60, 80 or 100) conservative substitutions.

- 10 Also encompassed by the invention are functional fragments of any of the above polypeptides.

- The polypeptides of the invention include fusion proteins containing a first domain and at least one additional domain. The first domain can be any of the B7-H3 or B7-H4 polypeptides described above or a functional fragment of any of these
- 15 polypeptides. The at least one additional domain can be a heterologous targeting or leader sequence, an amino acid sequence that facilitates purification, detection, or solubility of the fusion protein. The second domain can be, for example, all or part of an immunoglobulin (Ig) heavy chain constant region. Also included are isolated nucleic acid molecules encoding the fusion proteins.

- 20 The invention features vectors containing any of the DNAs of the invention and

- nucleic acid molecules encoding the fusion proteins of the invention. The vectors can be expression vectors in which the nucleic acid coding sequence or molecule is operably linked to a regulatory element which allows expression of the
- 25 nucleic acid sequence or molecule in a cell. Also included in the invention are cells (e.g., mammalian, insect, yeast, fungal, or bacterial cells) containing any of the vectors of the invention.

- Another embodiment of the invention is a method of co-stimulating a T cell that involves contacting the T cell with any of the B7-H3 or B7-H4 polypeptides of
- 30 the invention, functional fragments thereof, or fusion proteins of the invention; these 3 classes of molecule are, for convenience, designated "B7-H3 agents" or "B7-H4

agents." The contacting can be by, for example, culturing any of these B7-H3 or B7-H4 agents with the T cell *in vitro*. Alternatively, the T cell can be in a mammal and the contacting can be by, for example, administering any of the B7-H3 or B7-H4 agents to the mammal or administering a nucleic acid encoding the B7-H3 or B7-H4 agent to the mammal. In addition, the method can be an *ex vivo* procedure that involves: providing a recombinant cell which is the progeny of a cell obtained from the mammal and has been transfected or transformed *ex vivo* with a nucleic acid encoding any of the B7-H3 or B7-H4 agents so that the cell expresses the B7-H3 or B7-H4 agent; and administering the cell to the mammal. In this *ex vivo* procedure, the cell can be an antigen presenting cell (APC) that expresses the B7-H3 agent or B7-H4 agent on its surface. Furthermore, prior to administering to the mammal, the APC can be pulsed with an antigen or an antigenic peptide. In addition, the cell obtained from the mammal can be a tumor cell. In any of these methods of the invention, the B7-H3 or B7-H4 agents can co-stimulate the production of interferon- $\gamma$  by the T cell.

Also embodied by the invention is a method of co-stimulating a T cell in which the T cell is contacted with: (a) a first co-stimulatory polypeptide that can be either (i) B7-H1, (ii) B7-H2, (iii) B7-H3, (iv) B7-H4, (v) a functional fragment of any of (i) - (iv), or (vi) any of (i) - (v) but with one or more conservative substitutions; and (b) one or more additional co-stimulatory polypeptides that can be either (vi) B7-1, (vii) B7-2, (viii) B7-H1, (ix) B7-H2, (x) B7-H3, (xi) B7-H4, (xii) a functional fragment of any of (vi) - (xi), or (xii) any of (vi) - (xii) but with one or more conservative substitutions. The contacting can be by, for example, culturing the first co-stimulatory polypeptide and the one or more additional co-stimulatory polypeptides with the T cell *in vitro*. Alternatively, the T cell can be in a mammal and the contacting can be by, for example, administering the first co-stimulatory polypeptide and the one or more additional co-stimulatory polypeptides to the mammal. In addition, contacting of a T cell in a mammal can be by administering one or more nucleic acids encoding the first co-stimulatory polypeptide and the one or more additional co-stimulatory polypeptides to the mammal. The method can also be an *ex vivo* procedure that, for example, involves: providing a recombinant cell which is the progeny of a cell obtained from the mammal and which has been transfected or transformed *ex vivo* with one or more nucleic acids encoding the first co-stimulatory

polypeptide and the one or more additional polypeptides so that the cell expresses the first co-stimulatory polypeptide and the one or more additional co-stimulatory polypeptides; and administering the cell to the mammal. Alternatively, the *ex vivo* procedure can involve: providing a first recombinant cell which is the progeny of a cell obtained from the mammal and which has been transfected or transformed *ex vivo* with a nucleic acid encoding the first co-stimulatory polypeptide; providing one or more additional recombinant cells each of which is the progeny of a cell obtained from the mammal and each of which has been transfected or transformed *ex vivo* with a nucleic acid encoding one of the additional one or more co-stimulatory polypeptides; and administering the first cell and the one or more additional cells to the mammal. The recombinant cells used in the any of the *ex vivo* procedures can be antigen presenting cells (APC) and they can express the first co-stimulatory polypeptide and/or the one or more additional co-stimulatory polypeptides on their surfaces. Prior to the administering, APC can be pulsed with an antigen or an antigenic peptide. In addition, the cell obtained from the mammal can be a tumor cell.

In any of the above methods of co-stimulating a T cell, the mammal can be suspected of having, for example, an immunodeficiency disease, an inflammatory condition, or an autoimmune disease.

The invention includes a method of identifying a compound that inhibits an immune response. The method involves: providing a test compound; culturing, together, the compound, one or more B7-H3 or B7-H4 agents, a T cell, and a T cell activating stimulus; and determining whether the test compound inhibits the response of the T cell to the stimulus, as an indication that the test compound inhibits an immune response. The invention also embodies a method of identifying a compound that enhances an immune response. The method involves: providing a test compound; culturing, together, the compound, one or more of B7-H3 or B7-H4 agents, a T cell, and a T cell activating stimulus; and determining whether the test compound enhances the response of the T cell to the stimulus, as an indication that the test compound enhances an immune response. In both these methods, the stimulus can be, for example, an antibody that binds to a T cell receptor or a CD3 polypeptide. Alternatively, the stimulus can be an alloantigen or an antigenic peptide bound to a



major histocompatibility complex (MHC) molecule on the surface of an antigen presenting cell (APC). The APC can be transfected or transformed with a nucleic acid encoding the B7-H3 or B7-H4 agent and the B7-H3 or B7-H4 agent can be expressed on the surface of the APC.

5       The invention also features an antibody (e.g., a polyclonal or a monoclonal antibody) that binds specifically to one of the B7-H3 or B7-H4 polypeptides of the invention, e.g., the polypeptide with SEQ ID NO:1, SEQ ID NO:3, or SEQ ID NO:5.

10       The invention also features a method of producing any of the B7-H3 or B7-H4 polypeptides of the invention, functional fragments thereof, or fusion proteins of the invention. The method involves culturing a cell of the invention and purifying the relevant B7-H3 or B7-H4 protein from the culture.

15       "Polypeptide" and "protein" are used interchangeably and mean any peptide-linked chain of amino acids, regardless of length or post-translational modification. The invention also features B7-H3 and B7-H4 polypeptides with conservative substitutions. Conservative substitutions typically include substitutions within the following groups: glycine and alanine; valine, isoleucine, and leucine; aspartic acid and glutamic acid; asparagine, glutamine, serine and threonine; lysine, histidine and arginine; and phenylalanine and tyrosine.

20       The term "isolated" polypeptide or peptide fragment as used herein refers to a polypeptide or a peptide fragment which either has no naturally-occurring counterpart (e.g., a peptidomimetic), or has been separated or purified from components which naturally accompany it, e.g., in tissues such as pancreas, liver, spleen, ovary, testis, muscle, joint tissue, neural tissue, gastrointestinal tissue, or body fluids such as blood, serum, or urine. Typically, the polypeptide or peptide fragment is considered  
25       "isolated" when it is at least 70%, by dry weight, free from the proteins and naturally-occurring organic molecules with which it is naturally associated. Preferably, a preparation of a polypeptide (or peptide fragment thereof) of the invention is at least 80%, more preferably at least 90%, and most preferably at least 99%, by dry weight, the polypeptide (or the peptide fragment thereof), respectively, of the invention.  
30       Thus, for example, a preparation of polypeptide x is at least 80%, more preferably at least 90%, and most preferably at least 99%, by dry weight, polypeptide x. Since a

polypeptide that is chemically synthesized is, by its nature, separated from the components that naturally accompany it, the synthetic polypeptide or nucleic acid is "isolated."

An isolated polypeptide (or peptide fragment) of the invention can be  
5 obtained, for example, by extraction from a natural source (e.g., from human tissues or bodily fluids); by expression of a recombinant nucleic acid encoding the peptide; or by chemical synthesis. A peptide that is produced in a cellular system different from the source from which it naturally originates is "isolated," because it will be separated from components which naturally accompany it. The extent of isolation or purity can  
10 be measured by any appropriate method, e.g., column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis.

An "isolated DNA" means DNA free of the genes that flank the gene of interest in the genome of the organism in which the gene of interest naturally occurs. The term therefore includes a recombinant DNA incorporated into a vector, into an  
15 autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or eukaryote. It also includes a separate molecule such as: a cDNA where the corresponding genomic DNA has introns and therefore a different sequence; a genomic fragment; a fragment produced by polymerase chain reaction (PCR); a restriction fragment; a DNA encoding a non-naturally occurring protein, fusion  
20 protein, or fragment of a given protein; or a nucleic acid which is a degenerate variant of a naturally occurring nucleic acid. In addition, it includes a recombinant nucleotide sequence that is part of a hybrid gene, i.e., a gene encoding a fusion protein. Also included is a recombinant DNA that includes a portion of SEQ ID NO:2, SEQ ID NO:4, or SEQ ID NO:6. From the above it will be clear that an isolated DNA does  
25 not include a restriction fragment containing all or part of a gene that flanks the gene of interest in the genome of the organism in which the gene of interest naturally occurs. Furthermore, an isolated DNA does not mean a DNA present among hundreds to millions of other DNA molecules within, for example, cDNA or genomic DNA libraries or genomic DNA restriction digests in, for example, a restriction digest  
30 reaction mixture or an electrophoretic gel slice.

As used herein, a polypeptide that "co-stimulates" a T cell is a polypeptide that, upon interaction with a cell-surface molecule on the T cell, enhances the response of the T cell. The T cell response that results from the interaction will be greater than the response in the absence of the polypeptide. The response of the T cell in the absence of the co-stimulatory polypeptide can be no response or it can be a response significantly lower than in the presence of the co-stimulatory polypeptide. It is understood that the response of the T cell can be an effector, helper, or suppressive response.

As used herein, the term "co-stimulatory" polypeptide or molecule includes molecules such as B7-1, B7-2, B7-H1, B7-H2, B7-H3, B7-H4, 4-1BB, OX40, and herpes virus entry mediator (HVEM). As used herein, an "activating stimulus" is a molecule that delivers an activating signal to a T cell, preferably through the antigen specific T cell receptor (TCR). The activating stimulus can be sufficient to elicit a detectable response in the T cell. Alternatively, the T cell may require co-stimulation (e.g., by a B7-H3 or B7-H4 polypeptide) in order to respond detectably to the activating stimulus. Examples of activating stimuli include, without limitation, antibodies that bind to the TCR or to a polypeptide of the CD3 complex that is physically associated with the TCR on the T cell surface, alloantigens, or an antigenic peptide bound to a MHC molecule.

As used herein, a "fragment" of a B7-H3 or B7-H4 polypeptide is a fragment of the polypeptide that is shorter than the full-length polypeptide. Generally, fragments will be five or more amino acids in length. An antigenic fragment has the ability to be recognized and bound by an antibody.

As used herein, a "functional fragment" of a B7-H3 or B7-H4 polypeptide is a fragment of the polypeptide that is shorter than the full-length polypeptide and has the ability to co-stimulate a T cell. Methods of establishing whether a fragment of a B7-H3 or B7-H4 molecule is functional are known in the art. For example, fragments of interest can be made by either recombinant, synthetic, or proteolytic digestive methods. Such fragments can then be isolated and tested for their ability to co-stimulate T cells by procedures described herein.

As used herein, "operably linked" means incorporated into a genetic construct so that expression control sequences effectively control expression of a coding sequence of interest.

As used herein, the term "antibody" refers not only to whole antibody molecules, but also to antigen-binding fragments, e.g., Fab, F(ab')<sub>2</sub>, Fv, and single chain Fv fragments. Also included are chimeric antibodies.

As used herein, an antibody that "binds specifically" to an isolated B7-H4 polypeptide encoded by a DNA that includes a nucleic acid sequence that (i) encodes a polypeptide with the ability to co-stimulate a T cell and (ii) hybridizes under stringent conditions to the complement of a sequence that encodes a polypeptide with an amino acid sequence with SEQ ID NO:5, is an antibody that does not bind substantially to B7-1, B7-2, B7-H1, B7-H2, or B7-H3.

As used herein, an antibody that "binds specifically" to an isolated B7-H3 polypeptide that includes an amino acid sequence of amino acid residue 31 to amino acid residue 316 of SEQ ID NO: 3 is an antibody that does not bind substantially to B7-1, B7-2, B7-H1, B7-H2, or B7-H4. In addition, an antibody that binds specifically to B7-H3.1 preferably does not substantially bind to B7-H3.2 and *vice versa*.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. In case of conflict, the present document, including definitions, will control. Preferred methods and materials are described below, although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention. All publications, patent applications, patents and other references mentioned herein are incorporated by reference in their entirety. The materials, methods, and examples disclosed herein are illustrative only and not intended to be limiting.

Other features and advantages of the invention, e.g., enhancing immune responses in mammalian subjects, will be apparent from the following description, from the drawings and from the claims.

## BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a depiction of the nucleotide sequence of cDNA encoding B7H-3.1 (SEQ ID NO:2). Nucleotide residue 496 is shown in bold.

FIG. 2 is a depiction of the nucleotide sequence of cDNA encoding B7H-3.2 (SEQ ID NO:4). Nucleotide residue 496 is shown in bold.

FIG. 3 is a depiction of the amino acid sequence of B7-H3.1 (SEQ ID NO:1). Amino acid residue 166 is shown in bold.

FIG. 4 is a depiction of the amino acid sequence of B7-H3.2 (SEQ ID NO:3). Amino acid residue 166 is shown in bold.

FIG. 5A is a depiction of the amino acid sequences of human B7-H3.1 (SEQ ID NO:1), human B7-1 (SEQ ID NO:15), human B7-2 (SEQ ID NO:16), human B7-H1 (SEQ ID NO:17), and human B7-2 (SEQ ID NO:18) aligned for optimal homology. Identical amino acid residues are shaded in bold and conserved amino acid residues are boxed. Conserved cysteine residues are marked \*.

FIG. 5B is a depiction of the amino acid sequence of B7-H3.1 showing the predicted signal peptide (—), IgV-like domain (•••••), IgC-like domain (— — —), transmembrane domain (•••••), and intracellular tail (— • • — • •). Potential N-glycosylation sites are shown in bold ("N").

FIG. 6A is a pair of fluorescence flow cytometry histograms showing the staining by an anti-B7-H3 antiserum (open area) or control serum (shaded area) of 293 cells transfected with either the control pcDNA3.1(-) parental vector ("293/mock"; left histogram) or the pcDNA/B7-H3 vector containing the B7H-3.1 coding region ("293/B7-H3"; right histogram).

FIG. 6B is a series of fluorescence flow cytometry histograms showing the staining by an anti-B7-H3 antiserum (open area) or control serum (shaded area) of various cell types.

FIG. 7A is a series of fluorescence flow cytometry histograms showing the ability of B7-H3Ig fusion protein to bind to activated T cells but not to resting T cells. Unactivated nylon wool-purified T cells and nylon wool-purified T cells activated

with PHA (5 µg/ml) for 24 h, 48 h, or 72 h were stained with B7-H3Ig (5 µg; open areas) or control Ig (5 µg) prior to analysis.

FIG. 7B is a series of fluorescence flow cytometry histograms showing the ability of anti-B7-H3 antiserum, CTLA4Ig fusion protein, and ICOSIg fusion protein to bind to 293 cells transfected with expression vectors containing either a B7-H3.1 (solid line, shaded area), B7-1 (solid line, open area), or B7-H2 (dotted line, open area) coding sequence.

FIG. 8A is a line graph showing the proliferative response of nylon wool-purified T cells activated by anti-CD3 mAb (coated onto tissue culture well bottoms using a concentration of 40 ng/ml) and co-stimulated by either control Ig, B7-1Ig fusion protein, or B7-H3Ig fusion protein coated onto the tissue culture well bottoms at the indicated concentrations.

FIG. 8B is a bar graph showing the proliferative responses of purified CD4+ and CD8+ T cells activated by anti-CD3 mAb (coated onto tissue culture well bottoms using a concentration of 40 ng/ml) and co-stimulated by either control Ig, B7-1Ig fusion protein, or B7-H3Ig fusion protein, each coated onto tissue culture well bottoms using a concentration of 10 µg/ml.

FIG. 8C is a line graph showing the cytolytic activity on wild-type 624mel tumor target cells of cytotoxic T lymphocytes (CTL) (at the indicated effector to target cell ratios ("E:T")) generated by culturing nylon wool-purified T cells with 624mel tumor cells transfected with either the control parental pcDNA3.1(-) expression vector ("Mock/624mel") or the pcDNA/B7-H3 expression vector containing the B7H-3.1 coding region ("B7-H3/624mel").

FIG. 9 is a depiction of the nucleotide sequence of cDNA encoding B7-H4 (SEQ ID NO:6).

FIG. 10 is a depiction of the amino acid sequence of B7-H4 (SEQ ID NO:5).

FIG. 11 is a depiction of the amino acid sequences of the extracellular domains of human B7-H4 (SEQ ID NO: 19), human B7-H3.2 (SEQ ID NO:20), human B7-1 (SEQ ID NO:21), human B7-2 (SEQ ID NO:22), human B7-H1 (SEQ ID NO:23), and human B7-2 (SEQ ID NO:24) aligned for optimal homology. Identical

amino acid residues are shaded in bold and conserved amino acid residues are boxed. Conserved cysteine residues are marked \*.

FIG. 12 is a line graph showing the proliferative response ("CPM") of nylon wool-purified T cells activated by anti-CD3 mAb (coated onto tissue culture well bottoms using the indicated concentrations) and co-stimulated by either control human IgG ("hIgG"), B7-1Ig fusion protein, B7-H4Ig fusion protein, or B7-H4hIg coated onto tissue culture well bottoms at a concentration of 5 µg/ml.

FIG. 13 is a line graph showing the proliferative response ("CPM") of nylon wool-purified T cells activated by anti-CD3 mAb (coated onto tissue culture well bottoms using the indicated concentrations) and co-stimulated by either control human IgG ("hIgG"), B7-1Ig fusion protein, B7-H4Ig fusion protein, or B7-H4hIg fusion protein coated onto tissue culture well bottoms at a concentration of 10 µg/ml.

FIG. 14 is a bar graph showing the relative levels of mRNA encoding 23 cytokines in nylon wool purified T cells activated by anti-CD3 mAb (coated onto tissue culture well bottoms at a concentration of 500 ng/ml) and co-stimulated by either control human IgG (coated onto tissue culture well bottoms at a concentration of 5 µg/ml) ("hIgG"), B7-1Ig fusion protein (coated onto tissue culture well bottoms at a concentration of 5 µg/ml), or B7-H3Ig fusion protein (coated onto tissue culture well bottoms at a concentration of 5 µg/ml).

20

#### DETAILED DESCRIPTION

Using a novel PCR-based strategy, the inventor has identified two cDNA sequences (SEQ ID NOS:2 and 4) corresponding to two alleles of a gene encoding a novel B7-related molecule (B7-H3).

Translation of the cDNA sequences indicated that the two polypeptides (B7-H3.1 and B7-H3.2; SEQ ID NOS:1 and 3, respectively) encoded by the two allelic cDNA molecules are type I transmembrane proteins of 316 amino acids, each containing an immunoglobulin (Ig) V-like domain, Ig C-like domain, a transmembrane domain and a cytoplasmic domain of 30 amino acids. Northern blot analysis showed strong expression of the gene encoding B7-H3 in heart, liver,

30

placenta, prostate, testis, uterus, pancreas, small intestine, and colon, and weak expression in brain, skeletal muscle, kidney, and lung. Expression was undetectable in peripheral blood mononuclear cells (PBMC) but was detectable in spleen, lymph nodes, bone marrow, fetal liver, and thymus.

- 5           Fluorescence flow cytometry using an antiserum produced by immunization of mice with a carrier-conjugated peptide corresponding to a hydrophilic region of B7-H3, indicated no expression of B7-H3 on the majority of hematopoietic cells. However, less than 3% of resting CD14+ cells expressed the molecule on their surfaces. Activation with phorbol myristic acid (PMA) and ionomycin increased  
10           expression on T cells, monocytes, and dendritic cells.

Binding experiments indicated that T cells express a counter-receptor for B7-H3 that is not CTLA4, ICOS, or CD28.

- In vitro* experiments with isolated human T cells and a B7-H3.1-containing fusion protein indicated that while B7-H3 had no direct activity on T cells, it  
15           enhanced ("co-stimulated") CD4+ and CD8+ T cell proliferative responses induced by antibody specific for human CD3.

- Using a strategy similar to that used to clone B7-H3 cDNA, a cDNA molecule containing an open reading frame (orf) encoding another B7 homologue (B7-H4) was cloned, the nucleotide sequence of the orf (SEQ ID NO:6) was obtained, and the  
20           amino acid sequence of the encoded sequence (SEQ ID NO:5) was derived. B7-H4 is a type I transmembrane protein of 282 amino acids and has the same domain structure as B7-H3. Northern blot analysis showed expression of B7-H4 in lymphoid organs such as spleen and thymus. Binding experiments showed that there is a counter-receptor for B7-H4 on activated but not resting T cells. B7-H4, like B7-H3,  
25           co-stimulated T cell proliferation.

B7-H3 and B7-H4 can be used as augmenters of immune responses both *in vivo* and *in vitro*.



### Nucleic Acid Molecules

The B7-H3 and B7-H4 nucleic acid molecules of the invention can be cDNA, genomic DNA, synthetic DNA, or RNA, and can be double-stranded or single-stranded (*i.e.*, either a sense or an antisense strand). Fragments of these  
5 molecules are also considered within the scope of the invention, and can be produced by, for example, the polymerase chain reaction (PCR) or generated by treatment with one or more restriction endonucleases. A ribonucleic acid (RNA) molecule can be produced by *in vitro* transcription. Preferably, the nucleic acid molecules encode polypeptides that, regardless of length, are soluble under normal physiological  
10 conditions.

The nucleic acid molecules of the invention can contain naturally occurring sequences, or sequences that differ from those that occur naturally, but, due to the degeneracy of the genetic code, encode the same polypeptide (for example, the polypeptides with SEQ ID NOS:1, 3 or 5). In addition, these nucleic acid molecules  
15 are not limited to coding sequences, *e.g.*, they can include some or all of the non-coding sequences that lie upstream or downstream from a coding sequence.

The nucleic acid molecules of the invention can be synthesized (for example, by phosphoramidite-based synthesis) or obtained from a biological cell, such as the cell of a mammal. Thus, the nucleic acids can be those of a human, non-human  
20 primate (*e.g.*, monkey) mouse, rat, guinea pig, cow, sheep, horse, pig, rabbit, dog, or cat.

In addition, the isolated nucleic acid molecules of the invention encompass segments that are not found as such in the natural state. Thus, the invention encompasses recombinant nucleic acid molecules (for example, isolated nucleic acid  
25 molecules encoding B7-H3 or B7-H4) incorporated into a vector (for example, a plasmid or viral vector) or into the genome of a heterologous cell or into the genome of a homologous cell at a position other than the natural chromosomal location. Recombinant nucleic acid molecules and uses therefor are discussed further below.

Certain nucleic acid molecules of the invention are antisense molecules or are  
30 transcribed into antisense molecules. These can be used, for example, to down-regulate translation of B7-H3 or B7-H4 mRNA within a cell.

Techniques associated with detection or regulation of genes are well known to skilled artisans and such techniques can be used to diagnose and/or treat disorders associated with aberrant B7-H3 or B7-H4 expression. Nucleic acid molecules of the invention are discussed further below in the context of their therapeutic utility.

- 5 A B7-H3 or B7-H4 family gene or protein can be identified based on its similarity to the relevant B7-H3 or B7-H4 gene or protein, respectively. For example, the identification can be based on sequence identity. The invention features isolated nucleic acid molecules which are at least 50% (or 55%, 65%, 75%, 85%, 95%, or 98%) identical to: (a) a nucleic acid molecule that encodes the polypeptide of SEQ ID NO:1, 3, or 5; (b) the nucleotide sequence of SEQ ID NO:2, 4, or 6; or (c) a nucleic acid molecule which includes: (i) a segment of at least 30 (e.g., at least 50, 60, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 500, 550, 600, 650, 700, 800, 900, or 940) nucleotides of SEQ ID NO:2 or SEQ ID NO:4; or (ii) a segment of at least 30 (e.g., at least 50, 60, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 500, 550, 600, 650, 700, 800, or 840) nucleotides of SEQ ID NO:6.

- The determination of percent identity between two sequences is accomplished using the mathematical algorithm of Karlin and Altschul, *Proc. Natl. Acad. Sci. USA* 90, 5873-5877, 1993. Such an algorithm is incorporated into the BLASTN and BLASTP programs of Altschul et al. (1990) *J. Mol. Biol.* 215, 403-410. BLAST nucleotide searches are performed with the BLASTN program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to B7-H3- or B7-H4-encoding nucleic acids. BLAST protein searches are performed with the BLASTP program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to B7-H3 or B7-H4. To obtain gapped alignments for comparative purposes, Gapped BLAST is utilized as described in Altschul et al. (1997) *Nucleic Acids Res.* 25, 3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) are used (See <http://www.ncbi.nlm.nih.gov>).

- 30 Hybridization can also be used as a measure of homology between two nucleic acid sequences. A B7-H3- or B7-H4-encoding nucleic acid sequence, or a portion

thereof, can be used as hybridization probe according to standard hybridization techniques. The hybridization of a B7-H3 probe to DNA from a test source (e.g., a mammalian cell) is an indication of the presence of B7-H3 DNA in the test source and the hybridization of a B7-H4 probe to DNA from a test source (e.g., a mammalian cell) is an indication of the presence of B7-H4 DNA in the test source. Hybridization conditions are known to those skilled in the art and can be found in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y., 6.3.1-6.3.6, 1991. Moderate hybridization conditions are defined as equivalent to hybridization in 2X sodium chloride/sodium citrate (SSC) at 30°C, followed by one or more washes in 1 X SSC, 0.1% SDS at 50-60°C. Highly stringent conditions are defined as equivalent to hybridization in 6X sodium chloride/sodium citrate (SSC) at 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C.

The invention also encompasses: (a) vectors that contain any of the foregoing B7-H3- and B7-H4-related coding sequences and/or their complements (that is, "antisense" sequence); (b) expression vectors that contain any of the foregoing B7-H3-related and B7-H4-related coding sequences operatively associated with any transcriptional/translational regulatory elements (examples of which are given below) necessary to direct expression of the coding sequences; (c) expression vectors containing, in addition to sequences encoding a B7-H3 or a B7-H4 polypeptide, nucleic acid sequences that are unrelated to nucleic acid sequences encoding B7-H3 or a B7-H4, such as molecules encoding a reporter, marker, or a signal peptide, e.g., fused to B7-H3 or B7-H4; and (d) genetically engineered host cells that contain any of the foregoing expression vectors and thereby express the nucleic acid molecules of the invention.

Recombinant nucleic acid molecules can contain a sequence encoding B7-H3 or B7-H4, B7-H3 having a heterologous signal sequence, or B7-H4 having an heterologous signal sequence. The full length B7-H3 polypeptide, a domain of B7-H3, or a fragment thereof may be fused to additional polypeptides, as described below. In addition, the full-length B7-H4 polypeptide, a domain of B7-H4, or a fragment thereof may be fused to additional polypeptides, as described below. Similarly, the nucleic acid molecules of the invention can encode the mature form of

B7-H3 or B7-H4 or a form that includes an exogenous polypeptide which facilitates secretion.

The transcriptional/translational regulatory elements referred to above and which are further described below, include, but are not limited to, inducible and non-inducible promoters, enhancers, operators and other elements, which are known to those skilled in the art, and which drive or otherwise regulate gene expression. Such regulatory elements include but are not limited to the cytomegalovirus hCMV immediate early gene, the early or late promoters of SV40 adenovirus, the lac system, the trp system, the TAC system, the TRC system, the major operator and promoter regions of phage A, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase, the promoters of acid phosphatase, and the promoters of the yeast  $\alpha$ -mating factors.

Similarly, the nucleic acid can form part of a hybrid gene encoding additional polypeptide sequences, for example, sequences that function as a marker or reporter. Examples of marker or reporter genes include  $\beta$ -lactamase, chloramphenicol acetyltransferase (CAT), adenosine deaminase (ADA), aminoglycoside phosphotransferase (neo<sup>r</sup>, G418<sup>r</sup>), dihydrofolate reductase (DHFR), hygromycin-B-phosphotransferase (HPH), thymidine kinase (TK), lacZ (encoding  $\beta$ -galactosidase), and xanthine guanine phosphoribosyltransferase (XGPRT). As with many of the standard procedures associated with the practice of the invention, skilled artisans will be aware of additional useful reagents, for example, additional sequences that can serve the function of a marker or reporter. Generally, the hybrid polypeptide will include a first portion and a second portion; the first portion being a B7-H3 or B7-H4 polypeptide (or a fragment of such a polypeptide) and the second portion being, for example, the reporter described above or an Ig constant region or part of an Ig constant region, e.g., the CH2 and CH3 domains of IgG2a.

The expression systems that may be used for purposes of the invention include, but are not limited to, microorganisms such as bacteria (for example, *E. coli* and *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA, or cosmid DNA expression vectors containing the nucleic acid molecules of the invention; yeast (for example, *Saccharomyces* and *Pichia*) transformed with

recombinant yeast expression vectors containing the nucleic acid molecules of the invention, preferably containing the nucleic acid sequence encoding B7-H3 (SEQ ID NO:2 or 4) or B7-H4 (SEQ ID NO:6); insect cell systems infected with recombinant virus expression vectors (for example, baculovirus) containing the nucleic acid molecules of the invention; plant cell systems infected with recombinant virus expression vectors (for example, cauliflower mosaic virus (CaMV) and tobacco mosaic virus (TMV)) or transformed with recombinant plasmid expression vectors (for example, Ti plasmid) containing B7-H3 or B7-H4 nucleotide sequences; or mammalian cell systems (for example, COS, CHO, BHK, 293, VERO, HeLa, MDCK, WI38, and NIH 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (for example, the metallothionein promoter) or from mammalian viruses (for example, the adenovirus late promoter and the vaccinia virus 7.5K promoter). Also useful as host cells are primary or secondary cells obtained directly from a mammal transfected with a plasmid vector or infected with a viral vector.

#### Polypeptides and Polypeptide Fragments

The polypeptides of the invention include B7-H3, B7-H4, and functional fragments of these polypeptides. The polypeptides embraced by the invention also include fusion proteins which contain either full-length B7-H3 or B7-H4 or a functional fragment of either polypeptide fused to an unrelated amino acid sequence. The unrelated sequences can be additional functional domains or signal peptides. Signal peptides are described in greater detail and exemplified below. The polypeptides can also be any of those described above but with one or more conservative substitutions.

The polypeptides can be purified from natural sources (e.g., blood, serum plasma, tissues or cells such as T cells or any cell that naturally produces B7-H3 or B7-H4). Smaller peptides (less than 50 amino acids long) can also be conveniently synthesized by standard chemical means. In addition, both polypeptides and peptides can be produced by standard *in vitro* recombinant DNA techniques and *in vivo* recombination/genetic recombination (e.g., transgenesis), using the nucleotide

sequences encoding the appropriate polypeptides or peptides. Methods well known to those skilled in the art can be used to construct expression vectors containing relevant coding sequences and appropriate transcriptional/translational control signals. See, for example, the techniques described in Sambrook et al., *Molecular Cloning: A Laboratory Manual* (2nd Ed.) [Cold Spring Harbor Laboratory, N.Y., 1989], and Ausubel et al., *Current Protocols in Molecular Biology*, [Green Publishing Associates and Wiley Interscience, N.Y., 1989].

Polypeptides and fragments of the invention also include those described above, but modified for *in vivo* use by the addition, the amino- and/or carboxyl-terminal ends, of a blocking agent to facilitate survival of the relevant polypeptide *in vivo*. This can be useful in those situations in which the peptide termini tend to be degraded by proteases prior to cellular uptake. Such blocking agents can include, without limitation, additional related or unrelated peptide sequences that can be attached to the amino and/or carboxyl terminal residues of the peptide to be administered. This can be done either chemically during the synthesis of the peptide or by recombinant DNA technology by methods familiar to artisans of average skill.

Alternatively, blocking agents such as pyroglutamic acid or other molecules known in the art can be attached to the amino and/or carboxyl terminal residues, or the amino group at the amino terminus or carboxyl group at the carboxyl terminus can be replaced with a different moiety. Likewise, the peptides can be covalently or noncovalently coupled to pharmaceutically acceptable "carrier" proteins prior to administration.

Also of interest are peptidomimetic compounds that are designed based upon the amino acid sequences of the functional peptide fragments. Peptidomimetic compounds are synthetic compounds having a three-dimensional conformation (i.e., a "peptide motif") that is substantially the same as the three-dimensional conformation of a selected peptide. The peptide motif provides the peptidomimetic compound with the ability to co-stimulate T cells in a manner qualitatively identical to that of the B7-H3 or B7-H4 functional peptide fragment from which the peptidomimetic was derived. Peptidomimetic compounds can have additional characteristics that enhance

their therapeutic utility, such as increased cell permeability and prolonged biological half-life.

The peptidomimetics typically have a backbone that is partially or completely non-peptide, but with side groups that are identical to the side groups of the amino acid residues that occur in the peptide on which the peptidomimetic is based. Several  
5 types of chemical bonds, e.g., ester, thioester, thioamide, retroamide, reduced carbonyl, dimethylene and ketomethylene bonds, are known in the art to be generally useful substitutes for peptide bonds in the construction of protease-resistant peptidomimetics.

10

#### Methods of Co-stimulating a T Cell

The methods of the invention involve contacting a T cell with a B7-H3 or B7-H4 polypeptide of the invention, or a functional fragment thereof, in order to co-stimulate the T cell. Such polypeptides or functional fragments can have amino acid  
15 sequences identical to wild-type sequences or they can contain one or more conservative substitutions. The contacting can occur before, during, or after activation of the T cell. Contacting of the T cell with the B7-H3 or B7-H4 polypeptide will preferably be at substantially the same time as activation. Activation can be, for example, by exposing the T cell to an antibody that binds to the TCR or  
20 one of the polypeptides of the CD3 complex that is physically associated with the TCR. Alternatively, the T cell can be exposed to either an alloantigen (e.g., a MHC alloantigen) on, for example, an antigen presenting cell (APC) (e.g., a dendritic cell, a macrophage, a monocyte, or a B cell) or an antigenic peptide produced by processing of a protein antigen by any of the above APC and presented to the T cell by MHC  
25 molecules on the surface of the APC. The T cell can be a CD4+ T cell or a CD8+ T cell. The B7-H3 or B7-H4 polypeptide can be added to the solution containing the cells, or it can be expressed on the surface of an APC, e.g., an APC presenting an alloantigen or an antigen peptide bound to an MHC molecule. Alternatively, if the activation is *in vitro*, the B7-H3 or B7-H4 polypeptide can be bound to the floor of a  
30 the relevant culture vessel, e.g., a well of a plastic microtiter plate.

The methods can be performed *in vitro*, *in vivo*, or *ex vivo*. *In vitro* application of B7-H3 or B7-H4 can be useful, for example, in basic scientific studies of immune mechanisms or for production of activated T cells for use in either studies on T cell function or, for example, passive immunotherapy. Furthermore, B7-H3 or B7-H4 could be added to *in vitro* assays (e.g., in T cell proliferation assays) designed to test for immunity to an antigen of interest in a subject from which the T cells were obtained. Addition of B7-H3 or B7-H4 to such assays would be expected to result in a more potent, and therefore more readily detectable, *in vitro* response. However, the methods of the invention will preferably be *in vivo* or *ex vivo* (see below).

- 10 The B7-H3 and B7-H4 proteins and variants thereof are generally useful as immune response-stimulating therapeutics. For example, the polypeptides of the invention can be used for treatment of disease conditions characterized by immunosuppression: e.g., cancer, AIDS or AIDS-related complex, other virally or environmentally-induced conditions, and certain congenital immune deficiencies.
- 15 The polypeptides may also be employed to increase immune function that has been impaired by the use of radiotherapy of immunosuppressive drugs such as certain chemotherapeutic agents, and therefore are particularly useful when given in conjunction with such drugs or radiotherapy. The polypeptides can, furthermore, be used to enhance immune responses in normal subjects.
- 20 These methods of the invention can be applied to a wide range of species, e.g., humans, non-human primates, horses, cattle, pigs, sheep, goats, dogs, cats, rabbits, guinea pigs, hamsters, rats, and mice.

#### In Vivo Approaches

- 25 In one *in vivo* approach, a B7-H3 or B7-H4 polypeptide (or a functional fragment thereof) itself is administered to the subject. Generally, the compounds of the invention will be suspended in a pharmaceutically-acceptable carrier (e.g., physiological saline) and administered orally or by intravenous infusion, or injected subcutaneously, intramuscularly, intraperitoneally, intrarectally, intravaginally, intranasally, intragastrically, intratracheally, or intrapulmonarily. They are preferably delivered directly to an appropriate lymphoid tissue (e.g. spleen, lymph node, or
- 30



mucosal-associated lymphoid tissue (MALT)). The dosage required depends on the choice of the route of administration, the nature of the formulation, the nature of the patient's illness, the subject's size, weight, surface area, age, and sex, other drugs being administered, and the judgment of the attending physician. Suitable dosages are in the range of 0.01-100.0  $\mu\text{g/kg}$ . Wide variations in the needed dosage are to be expected in view of the variety of polypeptides and fragments available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by i.v. injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization as is well understood in the art. Administrations can be single or multiple (e.g., 2- or 3-, 4-, 6-, 8-, 10-, 20-, 50-, 100-, 150-, or more fold). Encapsulation of the polypeptide in a suitable delivery vehicle (e.g., polymeric microparticles or implantable devices) may increase the efficiency of delivery, particularly for oral delivery.

Alternatively, a polynucleotide containing a nucleic acid sequence encoding the B7-H3 or B7-H4 polypeptide or functional fragment thereof can be delivered to an appropriate cell of the animal. Expression of the coding sequence will preferably be directed to lymphoid tissue of the subject by, for example, delivery of the polynucleotide to the lymphoid tissue. This can be achieved by, for example, the use of a polymeric, biodegradable microparticle or microcapsule delivery vehicle, sized to optimize phagocytosis by phagocytic cells such as macrophages. For example, PLGA (poly-lacto-co-glycolide) microparticles approximately 1-10  $\mu\text{m}$  in diameter can be used. The polynucleotide is encapsulated in these microparticles, which are taken up by macrophages and gradually biodegraded within the cell, thereby releasing the polynucleotide. Once released, the DNA is expressed within the cell. A second type of microparticle is intended not to be taken up directly by cells, but rather to serve primarily as a slow-release reservoir of nucleic acid that is taken up by cells only upon release from the micro-particle through biodegradation. These polymeric particles should therefore be large enough to preclude phagocytosis i.e., larger than 5 $\mu\text{m}$  and preferably larger than 20 $\mu\text{m}$ .

Another way to achieve uptake of the nucleic acid is using liposomes, prepared by standard methods. The vectors can be incorporated alone into these delivery vehicles or co-incorporated with tissue-specific antibodies. Alternatively, one can prepare a molecular conjugate composed of a plasmid or other vector  
5 attached to poly-L-lysine by electrostatic or covalent forces. Poly-L-lysine binds to a ligand that can bind to a receptor on target cells [Cristiano et al. (1995), *J. Mol. Med.* 73, 479]. Alternatively, lymphoid tissue specific targeting can be achieved by the use of lymphoid tissue-specific transcriptional regulatory elements (TRE) such as a B lymphocyte, T lymphocyte, or dendritic cell specific TRE. Lymphoid tissue  
10 specific TRE are known [Thompson et al. (1992), *Mol. Cell. Biol.* 12, 1043-1053; Todd et al. (1993), *J. Exp. Med.* 177, 1663-1674; Penix et al. (1993), *J. Exp. Med.* 178, 1483-1496]. Delivery of "naked DNA" (i.e., without a delivery vehicle) to an intramuscular, intradermal, or subcutaneous site, is another means to achieve *in vivo* expression.

15 In the relevant polynucleotides (e.g., expression vectors) the nucleic acid sequence encoding the B7-H3 or B7-H4 polypeptide or functional fragment of interest with an initiator methionine and optionally a targeting sequence is operatively linked to a promoter or enhancer-promoter combination.

Short amino acid sequences can act as signals to direct proteins to specific  
20 intracellular compartments. For example, hydrophobic signal peptides (e.g., MAISGVPVLGFFILAVLMSAQESWA (SEQ ID NO:7)) are found at the amino terminus of proteins destined for the ER. While the sequence KFERQ (SEQ ID NO:8) (and other closely related sequences) is known to target intracellular polypeptides to lysosomes, other sequences (e.g., MDDQRDLISNNEQLP (SEQ ID  
25 NO:9) direct polypeptides to endosomes. In addition, the peptide sequence KDEL (SEQ ID NO:10) has been shown to act as a retention signal for the ER. Each of these signal peptides, or a combination thereof, can be used to traffic the B7-H3 or B7-H4 polypeptides or functional fragments of the invention as desired. DNAs encoding the B7-H3 or B7-H4 polypeptides or functional fragments containing  
30 targeting signals will be generated by PCR or other standard genetic engineering or synthetic techniques.

A promoter is a TRE composed of a region of a DNA molecule, typically within 100 basepairs upstream of the point at which transcription starts. Enhancers provide expression specificity in terms of time, location, and level. Unlike a promoter, an enhancer can function when located at variable distances from the transcription site, provided a promoter is present. An enhancer can also be located downstream of the transcription initiation site. To bring a coding sequence under the control of a promoter, it is necessary to position the translation initiation site of the translational reading frame of the peptide or polypeptide between one and about fifty nucleotides downstream (3') of the promoter. The coding sequence of the expression vector is operatively linked to a transcription terminating region.

Suitable expression vectors include plasmids and viral vectors such as herpes viruses, retroviruses, vaccinia viruses, attenuated vaccinia viruses, canary pox viruses, adenoviruses and adeno-associated viruses, among others.

Polynucleotides can be administered in a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers are biologically compatible vehicles which are suitable for administration to a human, e.g., physiological saline. A therapeutically effective amount is an amount of the polynucleotide which is capable of producing a medically desirable result (e.g., an enhanced T cell response) in a treated animal. As is well known in the medical arts, the dosage for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. Dosages will vary, but a preferred dosage for administration of polynucleotide is from approximately  $10^6$  to  $10^{12}$  copies of the polynucleotide molecule. This dose can be repeatedly administered, as needed. Routes of administration can be any of those listed above.

Included in these *in vivo* approaches, are methods of co-stimulating a T cell that involve administering more than one co-stimulatory molecule or functional fragment thereof. Such combinations can be any combination of one or more of co-stimulatory polypeptides, e.g., B7-1, B7-2, B7-H1, B7-H2, B7-H3, B7-H4, 4-1BB, OX40, or HVEM and functional fragments of any of these. The proteins or functional fragments *per se* can be administered (as above) or nucleic acids (e.g., expression

vectors) encoding the proteins or functional fragments can be administered (as above). Where expression vectors are used, a single vector containing coding sequences for two or more of the co-stimulatory polypeptides or functional fragments can be administered. Alternatively, multiple (e.g., 2, 3, 4, 5, or 6) individual vectors, each encoding one or more (e.g., 2, 3, 4, 5, or 6) of the co-stimulatory polypeptides or functional fragments thereof can be administered.

#### Ex Vivo Approaches

Peripheral blood mononuclear cells (PBMC) can be withdrawn from the patient or a suitable donor and exposed *ex vivo* to an activating stimulus (see above) and a B7-H3 or B7-H4 polypeptide or polypeptide fragment (whether in soluble form or attached to a solid support by standard methodologies). The PBMC containing highly activated T cells are then introduced into the same or a different patient.

An alternative *ex vivo* strategy can involve transfecting or transducing cells obtained from the subject with a polynucleotide encoding a B7-H3 or B7-H4 polypeptide or functional fragment-encoding nucleic acid sequences described above. The transfected or transduced cells are then returned to the subject. While such cells would preferably be hemopoietic cells (e.g., bone marrow cells, macrophages, monocytes, dendritic cells, or B cells) they could also be any of a wide range of types including, without limitation, fibroblasts, epithelial cells, endothelial cells, keratinocytes, or muscle cells in which they act as a source of the B7-H3 or B7-H4 polypeptide or functional fragments thereof for as long as they survive in the subject. The use of hemopoietic cells, that include the above APC, would be particularly advantageous in that such cells would be expected to home to, among others, lymphoid tissue (e.g., lymph nodes or spleen) and thus the B7-H3 or B7-H4 polypeptide or functional fragment would be produced in high concentration at the site where they exert their effect, i.e., enhancement of an immune response. In addition, if APC are used, the APC expressing the exogenous B7-H3 or B7-H4 molecule can be the same APC that presents an alloantigen or antigenic peptide to the relevant T cell. The B7-H3 or B7-H4 polypeptides can be secreted by the APC or expressed on its surface. Prior to returning the recombinant APC to the patient, they

can optionally be exposed to sources of antigens or antigenic peptides of interest, e.g., those of tumors, infectious microorganisms, or autoantigens. The same genetic constructs and trafficking sequences described for the *in vivo* approach can be used for this *ex vivo* strategy. Furthermore, tumor cells, preferably obtained from a patient, can be transfected or transformed by a vector encoding a B7-H3 or B7-H4 polypeptide or functional fragment thereof. The tumor cells, preferably treated with an agent (e.g., ionizing irradiation) that ablates their proliferative capacity, are then returned to the patient where, due to their expression of the exogenous B7-H3 or B7-H4 (on their cell surface or by secretion), they can stimulate enhanced tumoricidal T cell immune responses. It is understood that the tumor cells which, after transfection or transformation, are injected into the patient, can also have been originally obtained from an individual other than the patient.

The *ex vivo* methods include the steps of harvesting cells from a subject, culturing the cells, transducing them with an expression vector, and maintaining the cells under conditions suitable for expression of the B7-H3 or B7-H4 polypeptide or functional fragment. These methods are known in the art of molecular biology. The transduction step is accomplished by any standard means used for *ex vivo* gene therapy, including calcium phosphate, lipofection, electroporation, viral infection, and biolistic gene transfer. Alternatively, liposomes or polymeric microparticles can be used. Cells that have been successfully transduced are then selected, for example, for expression of the coding sequence or of a drug resistance gene. The cells may then be lethally irradiated (if desired) and injected or implanted into the patient.

It is understood that in these *ex vivo* procedures, the cells to be introduced into a subject can be transfected or transformed with one or more (e.g., two, three, four, five, or six) expression vectors containing one or more (e.g., two, three, four, five, or six) sequences encoding any of the co-stimulatory molecules listed above (e.g., B7-1, B7-2, B7-H1, B7-H2, B7-H3, or B7-H4) or functional fragments thereof prior to introduction.

Methods of Screening for Compounds that Inhibit or Enhance Immune Responses.

The invention provides methods for testing compounds (small molecules or macromolecules) that inhibit or enhance an immune response. Such a method can involve, e.g., culturing a B7-H3 or B7-H4 polypeptide of the invention (or a  
5 functional fragment thereof) with T cells in the presence of a T cell stimulus (see above). The B7-H3 or B7-H4 molecule can be in solution or membrane bound (e.g., expressed on the surface of the T cells) and it can be natural or recombinant. Furthermore, the B7-H3 or B7-H4 polypeptides (or functional fragments thereof) can have amino acid sequences identical to wild-type sequences or they can have one or  
10 more conservative substitutions. Compounds that inhibit the T cell response will likely be compounds that inhibit an immune response while those that enhance the T cell response will likely be compounds that enhance an immune response.

The invention also relates to using B7-H3 or B7-H4 or functional fragments thereof to screen for immunomodulatory compounds that can interact with B7-H3 or  
15 B7-H4. One of skill in the art would know how to use standard molecular modeling or other techniques to identify small molecules that would bind to T cell interactive sites of B7-H3 or B7-H4. One such example is provided in Broughton (1997) Curr. Opin. Chem. Biol. 1, 392-398.

A candidate compound whose presence requires at least 1.5-fold (e.g., 2-fold, 4-fold, 6-fold, 10-fold, 150-fold, 1000-fold, 10,000-fold, or 100,000-fold) more B7-  
20 H3 or B7-H4 in order to achieve a defined arbitrary level of T cell activation than in the absence of the compound can be useful for inhibiting an immune response. On the other hand, a candidate compound whose presence requires at least 1.5 fold (e.g., 2-fold, 4-fold, 6-fold, 10-fold, 100-fold, 1000-fold, 10,000 fold, or 100,000-fold) less  
25 B7-H3 or B7-H4 to achieve a defined arbitrary level of T cell activation than in the absence of the compound can be useful for enhancing an immune response. Compounds capable of interfering with or modulating B7-H3 or B7-H4 function are good candidates for immunosuppressive immunoregulatory agents, e.g., to modulate an autoimmune response or suppress allogeneic or xenogeneic graft rejection.

30

### B7-H3 and B7-H4 Antibodies

The invention features antibodies that bind to the B7-H3 or B7-H4 polypeptides or fragments of such polypeptides. Such antibodies can be polyclonal antibodies present in the serum or plasma of animals (e.g., mice, rabbits, rats, guinea pigs, sheep, horses, goats, cows, or pigs) which have been immunized with the relevant B7-H3 or B7-H4 polypeptide or peptide fragment using methods, and optionally adjuvants, known in the art. Such polyclonal antibodies can be isolated from serum or plasma by methods known in the art. Monoclonal antibodies that bind to the above polypeptides or fragments are also embodied by the invention. Methods of making and screening monoclonal antibodies are well known in the art.

Once the desired antibody-producing hybridoma has been selected and cloned, the resultant antibody can be produced in a number of methods known in the art. For example, the hybridoma can be cultured *in vitro* in a suitable medium for a suitable length of time, followed by the recovery of the desired antibody from the supernatant. The length of time and medium are known or can be readily determined.

Additionally, recombinant antibodies specific for B7-H3 or B7-H4, such as chimeric and humanized monoclonal antibodies comprising both human and non-human portions, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example, using methods described in Robinson et al., International Patent Publication PCT/US86/02269; Akira et al., European Patent Application 184,187; Taniguchi, European Patent Application 171,496; Morrison et al., European Patent Application 173,494; Neuberger et al., PCT Application WO 86/01533; Cabilly et al., U.S. Patent No. 4,816,567; Cabilly et al., European Patent Application 125,023; Better et al. (1988) *Science* 240, 1041-43; Liu et al. (1987) *J. Immunol.* 139, 3521-26; Sun et al. (1987) *PNAS* 84, 214-18; Nishimura et al. (1987) *Canc. Res.* 47, 999-1005; Wood et al. (1985) *Nature* 314, 446-49; Shaw et al. (1988) *J. Natl. Cancer Inst.* 80, 1553-59; Morrison, (1985) *Science* 229, 1202-07; Oi et al. (1986) *BioTechniques* 4, 214; Winter, U.S. Patent No. 5,225,539; Jones et al. (1986) *Nature* 321, 552-25; Veroeyan et al. (1988) *Science* 239, 1534; and Beidler et al. (1988) *J. Immunol.* 141, 4053-60.

Also included within the scope of the invention are antibody fragments and derivatives which contain at least the functional portion of the antigen binding domain of an antibody that binds specifically to B7-H3 or B7-H4. Antibody fragments that contain the binding domain of the molecule can be generated by known techniques.

- 5 For example, such fragments include, but are not limited to:  $F(ab')_2$  fragments which can be produced by pepsin digestion of antibody molecules; Fab fragments which can be generated by reducing the disulfide bridges of  $F(ab')_2$  fragments; and Fab fragments which can be generated by treating antibody molecules with papain and a reducing agent. See, e.g., National Institutes of Health, 1 Current Protocols In  
10 Immunology, Coligan *et al.*, ed. 2.8, 2.10 (Wiley Interscience, 1991). Antibody fragments also include Fv (e.g., single chain Fv (scFv)) fragments, *i.e.*, antibody products in which there are no constant region amino acid residues. Such fragments can be produced, for example, as described in U.S. Patent No. 4,642,334 which is incorporated herein by reference in its entirety.

- 15 The following examples are meant to illustrate, not limit, the invention.

#### Example 1. Materials and Methods

- Cell lines, cell preparation and activation. The adenovirus-transformed human kidney epithelial line 293 and the histocytic lymphoma line U937 were purchased from ATCC (Manassas, VA). The 624mel melanoma line was a gift from Dr. Rong-  
20 Fu Wang (Surgery Branch, NCI). Cell lines were grown and maintained in DMEM or RPMI 1640 media (GIBCO, Gaithersburg, MD) supplemented with 10% fetal bovine serum (FBS) (HyClone, Logan, UT), 25mM HEPES (Life Technologies, Grand Island, USA), 2 mM L-glutamine (Life Technologies), 100 U/ml penicillin (Life Technologies), 100  $\mu$ g/ml streptomycin (Life Technologies), and  $5 \times 10^{-5}$  M 2-  
25 mercaptoethanol (VWR, Chicago, IL). Methods for the preparation of peripheral blood mononuclear cells (PBMC) from healthy donors and monocyte-derived dendritic cells (DC) were described previously [Chapoval *et al.* (2000) *Blood*. 95, 2346-2351]. Fluorescence flow cytometry (FFC) analysis using monoclonal antibodies (mAb) specific for B7-1, B7-2, and MHC class II indicated that >99% of  
30 cells in the monocyte-derived DC preparations had a typical DC phenotype. Total T cell-enriched populations were obtained by passing nonadherent PBMC through



nylon wool columns as described previously [Tamada et al. (2000) *J. Immunol.* 164, 4105-4110]. CD4<sup>+</sup> and CD8<sup>+</sup> T cell subsets were further purified using the MACS magnetic bead system (Miltenyl Biotec, Auburn, CA) according to the manufacturer's instructions. The purity of the T cell subset preparations were >98% based on FFC analysis using T cell subset-specific mAb. For the activation of U937 cells or DC, the cells ( $1 \times 10^6$  cells/ml) were cultured for 24 h with lipopolysaccharide (LPS; 1 µg/ml). Adherent PBMC were activated with a combination of LPS (100 ng/ml) and human interferon-γ (IFN-γ; 1500 IU/ml; R&D Systems, Minneapolis MN). For activation of T and B cells, PBMC ( $2.5 \times 10^6$  cells/ml) were cultured for 24 h with phytohemagglutinin (PHA; 5 µg/ml) or LPS (1 µg/ml), respectively. All cell types were also activated with phorbol myristic acid (PMA; 5 ng/ml) and ionomycin (250 ng/ml).

Cloning and sequencing of full-length human B7-H3 cDNA. The National Center for Biotechnology Information (NCBI) and Human Genome Sciences, Inc. expressed sequence tag (EST) databases containing the sequences of >500 different cDNA libraries were screened for cDNA sequences having homology to published B7-1, B7-2, B7-H1, and B7-H2 sequences using the BLASTN and TBLASTN algorithms. Information obtained from this screening was used to generate appropriate PCR primers and full-length human cDNA molecules encoding B7-H3 was generated by PCR from a THP-1 cDNA library prepared by the SMART PCR cDNA synthesis kit (Clontech, Palo Alto, CA). The resulting PCR product was cloned into the pcDNA3.1(-) vector (Invitrogen, Carlsbad, CA). The nucleotide sequence of the B7-H3 encoding cDNA molecules was verified by sequencing of several independent clones. For transient expression of the B7-H3 gene, the vector containing full-length B7-H3.1 encoding cDNA (pcDNA/B7-H3) or control parental vector (pcDNA3.1(-)) was transfected into 293 or 624mel cells by Fugene 6 (Boehringer-Mannheim) according to the manufacturer's instructions.

Production of fusion proteins. Recombinant B7-H3Ig fusion protein was prepared by fusing the coding region of the extracellular domain of B7-H3 to the Fc constant region of mouse IgG2a as described previously [Chapoval et al. (2000). *Methods Mol. Med.* 45:247-255]. An expression vector containing a DNA sequence

encoding B7-H3Ig was transfected into 293 cells by calcium phosphate precipitation and cultured in serum-free DMEM. The supernatant was collected at 72 h and the fusion protein was purified by Protein G sepharose columns (Pharmacia, Uppsala, Sweden). The purity and expected molecular weight of the fusion protein were confirmed by electrophoresis on polyacrylamide gels and by Western blot using the anti-B7-H3 antibody described below. Analogous fusion proteins (ICOSIg, B7-1Ig, and CTLA4Ig) containing the extracellular domains of ICOS, B7-1, and CTLA4, respectively, were prepared by a similar method.

RNA analysis. The expression of B7-H3 mRNA in human tissues was analyzed using human multiple-tissue and tumor line Northern blots purchased from Clontech (Palo Alto, CA). Full-length cDNA encoding B7-H3 was labeled with [<sup>32</sup>P]-dCTP and hybridized to filter membranes according to the manufacturer's protocol.

Immunization, antibody production and FACS analysis. For production of antibody specific for human B7-H3, a peptide consisting of amino acids 142-165 (YSKPSMTLEPNKDLRPGDTVTITC) (SEQ ID NO:12) spanning a hydrophilic region of B7-H3 was synthesized and conjugated to KLH as described previously [Tamada et al. (2000) *J.Immunol.* 164, 4105-4110]. Polyclonal antibodies were prepared by immunizing BALB/c mice with the peptide in complete Freund's adjuvant (Sigma) and subsequently boosting two times with the peptides in incomplete Freund's adjuvant. Blood was collected and sera prepared 10 days after the final boost. The specificity of the antiserum was determined by ELISA against B7-H3Ig and indirect immunofluorescence of 293 cells transfected to express B7-H3, B7-H2, B7-H1, and B7-1. Serum prepared from the blood of non-immunized BALB/c mice was used as a negative control. Substantially identical results were obtained with two antisera generated in the same way by immunization of mice with B7-H3 hydrophilic region peptides corresponding to amino acid residues 166-189 (SSYRGYPEAEVFWQDGQGVPLTGN) (SEQ ID NO:13) of B7-H3.1 or amino acid residues 223-247 (RNPVLQQDAHGSVTTTGQPMTFPPE) (SEQ ID NO:14) of B7-H3.

For analysis of B7-H3 expression and B7-H3 interaction with a putative counter-receptor, cells were incubated either with anti-B7-H3 antibodies (1:1000

dilution in PBS), CTLA4Ig (5 µg/sample) or ICOSIg (5 µg/sample) on ice. After a 45 min incubation, cells were washed and cultured for 45 min with fluorescein isothiocyanate (FITC)-conjugated goat anti-mouse Ig F(ab')<sub>2</sub> (BioSource, Camarillo, CA). For two color staining, cells were washed and further incubated with phycoerythrin (PE)-conjugated mAb specific for CD3, CD14, or CD19 (PharMingen, San Diego, CA). For the analysis of CD3+, CD19+, and CD14+ cells, peripheral blood mononuclear cells (PBMC) (untreated or activated as described below) were stained with anti-B3-H7 antiserum (green fluorescence) and either anti-CD3, anti-CD19, or anti-CD14 monoclonal antibody (mAb) (orange fluorescence). The proportion of B7-H3 expressing cells (green fluorescence) within the CD3+, the CD19+, and the CD14+ populations was assessed by gating the flow cytometer on the relevant populations (orange fluorescence). Expression of a B7-H3 counter-receptor was analyzed by incubation of cells with B7-H3Ig (5 µg/sample) with subsequent staining using FITC-conjugated goat anti-mouse Ig F(ab')<sub>2</sub>. Single or double stained cells were analyzed using the Becton Dickinson FACScan (Mountain View, CA).

T cell proliferation and cytokine production assay. T cell proliferation was measured as previously described [Dong et al. (1999) *Nat. Med.* 5, 1365-1369]. Briefly, flat-bottom 96-well microtiter culture plates were coated at 4°C overnight with 50 µl /well of anti-CD3 mAb (40 ng/ml) and subsequently coated with the indicated concentrations of B7-H3Ig, B7-1Ig, or control Ig at 37°C for 4 h. T cells were added to the wells at the indicated concentrations. For measurements of T cell proliferation, the plates were cultured for 72 h and [<sup>3</sup>H]-thymidine (1 µCi/well) was added 18 h prior to harvesting of the cultures. [<sup>3</sup>H]-thymidine incorporation was measured with a MicroBeta Trilix liquid scintillation counter (Wallac, Turku, Finland). For measurements of cytokine (interleukin- (IL-)2, IL-10, and interferon-γ (IFN-γ) production, supernatants were removed from the cultures at the indicated times after culture initiation and subjected to ELISA for the cytokines using commercially available ELISA kits (PharMingen).

CTL generation and cytotoxicity assay. CTL were generated in 24-well culture plates by co-culturing of nylon wool column-purified T cells (5 × 10<sup>6</sup> cells/well) in the presence of irradiated (10,000 rad) 624mel cells (1 × 10<sup>5</sup> cells/well)

transfected with either pcDNA/B7-H3 or control parent pcDNA3.1(-) plasmid. The cultured cells were harvested on day 5 and CTL activity against parental 624mel was measured in a standard 4h  $^{51}\text{Cr}$ -release assay [Chapoval et al. (1998) *J.Immunol.* 161, 6977-6984].

- 5        Cytokine gene expression. Total RNA was prepared using TRI Reagent (Sigma) from  $5 \times 10^6$  T cells which had been cultured in 24-well tissue culture plates for 72h with a suboptimal dose of anti-CD3 antibody (coated onto tissue culture plate wells at a concentration of 200 ng/ml) and with either control Ig (coated onto tissue culture plate wells at a concentration of 5 ug/ml), B7-1Ig (coated onto tissue culture plate wells at a concentration of 5 ug/ml), or B7-H3Ig (coated onto tissue culture plate wells at a concentration of 5 ug/ml). Ten micrograms of the total RNA from each of the three cell populations were used as a template to produce  $^{32}\text{P}$  labeled cDNA by standard procedures using retroviral (MMTV) reverse transcriptase (Promega, Madison, WI) and [ $\alpha$ - $^{32}\text{P}$ ]-dCTP (NEN, Boston, MA). The Human Common
- 10        Cytokine-1 GEAarray Kit (No hGEA9912099, SuperArray Inc. Bethesda, MD) was used to determine the relative amounts of cDNA encoding 23 different cytokines (see Fig. 14) in the three  $^{32}\text{P}$  labeled cDNA samples as an indirect measure of the relative amounts of corresponding mRNA in the three cell populations. The kit includes cytokine mRNA arrays which are membranes containing mRNA encoding
- 15        the 23 cytokines and two house keeping protein, beta-actin and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) bound to discrete areas of the membrane. A separate array was exposed under standard hybridization conditions to each of the three  $^{32}\text{P}$  labeled cDNA samples and the three arrays were washed under washing conditions specified by the manufacturer. A STORM Phosphoimager system
- 20        (Molecular Dynamics, Sunnyvale, CA) was used to measure the signals due to hybridization of  $^{32}\text{P}$  labeled cDNA to appropriate areas on the membranes. The data are expressed as arbitrary units calculated by the following formula:
- 25

$$\text{Arbitrary units} = (\text{Cytokine signal} - \text{background signal}) / (\text{beta-actin signal} - \text{background signal})$$

### Example 2. Molecular Cloning and Expression Pattern of B7-H3-encoding DNA

The EST databases of NCBI and Human Genome Sciences, Inc. were screened for sequences homologous to the DNA sequences encoding the extracellular regions of all published B7 family members. By a PCR-based strategy using primers based on EST sequences of interest, two open reading frames (orf) encoding two different allelic forms of a B7-like molecule were identified in a human dendritic cell (DC)-derived cDNA library. The nucleic acid sequences were confirmed by analyses of independent RT-PCR products from human THP-1 and DC library. The two sequences, which differed by a single nucleotide and most likely corresponding to B7-H3 alleles, were observed by sequencing several independent clones obtained from several independent PCRs. In one allele (SEQ ID NO:2) (B7H-3.1) nucleotide 496 is a T (Fig. 1) and in the second allele (B7H-3.2) (SEQ ID NO:4) nucleotide 496 is a C (Fig. 2). At the protein level, B7-H3.1 (SEQ ID NO:1) contains a serine residue at position 166 (Fig. 3) and B7-H3.2 (SEQ ID NO: 2) contains a proline residue at position 166 (Fig. 4). Except at position 166, B7-H3.1 and B7-H3.2 have identical amino acid sequences. The B7-H3.1 orf encodes a putative 316 amino acid protein (B7-H3) that shares identity in its predicted extracellular receptor-binding domain with human B7-H1 (27%), B7-H2 (25%), B7-1 (21%), and B7-2 (20%) (Fig 5A). The putative B7-H3 protein contains a typical signal peptide in its N-terminus, a single extracellular V-like Ig domain and a single C-like Ig domains, a transmembrane region, and a 45 amino acid cytoplasmic tail (Fig. 5B), indicating that B7-H3 is a type I transmembrane protein belonging to the Ig superfamily. Similar to other members of the family, B7-H3 has four conserved cysteine residues that are believed to be involved in the formation of V- and C-like Ig domains. The absence of a heptad structure and B30.2 domains makes B7-H3 distinct from butyrophilins and myelin oligodendrocyte glycoproteins [reviewed in Linsley, et al. (1994) *Protein Sci.* 3, 1341-1343; Henry et al. (1999) *Immunol.Today* 20, 285-288]. The above data indicate that B7-H3 is a member of the B7 costimulatory ligand family.

Northern blot analysis indicated that B7-H3 is encoded by a single 4.1-kb mRNA and is expressed at high levels in many normal human tissues including heart, liver, placenta, prostate, testis, uterus, pancreas, small intestine, and colon; low levels

of B7-H3 mRNA were also found in brain, skeletal muscle, kidney, and lung. B7-H3 mRNA could also be detected in several lymphoid organs including spleen, lymph nodes, bone marrow, fetal liver, and thymus. Surprisingly, no B7-H3 mRNA was detected in RNA from PBMC in two independent northern blots. Several tumor lines, including melanoma G361, cervix adenocarcinoma HeLa S3, chronic myelogenous leukemia K562, lung carcinoma A546, and colorectal adenocarcinoma SW480, also express B7-H3 mRNA. Molt-4 (lymphoblastic leukemia) and Raji (Burkitts lymphoma) were negative for B7-H3 mRNA. Equal levels of actin mRNA were observed in all the blots. Using RT-PCR, B7-H3 mRNA was detectable in RNA from K562, U937, THP-1, and dendritic cells but not in Raji, Jurkat, Molt-4, or a T cell clone.

FFC analysis was performed to detect cell-surface expression of B7-H3 protein. Polyclonal antibodies specific for human B7-H3 were generated by immunizing mice with KLH-conjugated synthetic peptides spanning the hydrophilic regions of human B7-H3. Results obtained using an antiserum generated by immunization with a peptide containing amino acid residues 142-165 are described. The antiserum stained 293 cells transiently transfected to express B7-H3, but not B7-H1, B7-H2, nor B7-1 (Fig. 6A, 7B and data not shown), indicating that this antiserum is specific for B7-H3 protein. B7-H3 is detectable on a small fraction of resting CD14<sup>+</sup> cells (<3%), but not on resting CD3<sup>+</sup> and CD19<sup>+</sup> cells (Fig. 6B). Activation ("standard activation") of PBMC by various methods including PHA, LPS, or a combination of LPS and IFN- $\gamma$  did not modulate B7-H3 expression. In contrast, stimulation with a combination of PMA and ionomycin significantly increased surface expression of B7-H3 on CD3<sup>+</sup> and CD14<sup>+</sup> cells but not on CD19<sup>+</sup> cells (Fig. 6B). Interestingly, treatment by PMA and ionomycin (but not by LPS) also increased the expression of B7-H3 on cytokine-induced DC and on cells of the human monocytic tumor line U937 (Fig. 6B). The epithelium-derived tumor lines including choriocarcinoma BeWo, colorectal adenocarcinomas HT29, WiDr, and SW620 negative for B7-H3, with or without activation by PMA and ionomycin (data not shown). These data indicate that surface B7-H3 is not constitutively expressed on the majority of hematopoietic cells but is selectively induced by PMA and ionomycin in T cells, monocytes, and DC.

### Example 3. Expression of a B7-H3 Counter-Receptor

To determine whether a counter-receptor (B7-H3CR) for B7-H3 is expressed on T cells, a B7-H3Ig fusion protein was prepared by fusing the extracellular domain of B7-H3 and the Fc portion of mouse IgG2a. FFC analysis (Fig. 7A) indicated that  
5 purified resting T cells do not express a B7-H3CR. Stimulation of T cells with PHA, however, led to a rapid up-regulation of B7-H3CR within the first 24 h of culture. The expression waned after 48 h (Fig 7A).

293 cells were transiently transfected with pcDNA/B7-H3, stained with  
10 CTLA4Ig and ICOSIg, and subjected to FACS analysis. The cells were also stained with anti-B7-H3 antibodies as a positive control. Neither CTLA4Ig nor ICOSIg stained B7-H3/293 cells (Fig. 7B). Furthermore, while anti-B7-H3 antibodies did not bind to B7-1/293 or B7-H2/293, CTLA4Ig bound to B7-1/293 and ICOSIg bound to B7-H2/293 (Fig. 7B). Thus B7-H3 is a ligand for an inducible T cell counter-receptor  
15 distinct from CTLA-4 and ICOS.

### Example 4. B7-H3 Co-stimulates T Cell Proliferation and the Generation of CTL

A previously described co-stimulation assay was used [Dong et al. (1999) *Nat.Med.* 5, 1365-1369] to test whether B7-H3 co-stimulates the proliferation of T  
20 cells. In this assay, purified T cells were stimulated by immobilized anti-CD3 mAb in the presence of immobilized B7-H3Ig. The proliferation of T cells was determined by [<sup>3</sup>H]-thymidine-incorporation after a 72 h incubation. B7-H3Ig increased T cell proliferation in a dose-dependent fashion (Fig. 8A) in the presence of a suboptimal dose of anti-CD3 mAb (coated onto the plates at a concentration of 40 ng/ml).  
25 Interestingly, B7-1Ig coated onto the plates at concentrations of 2.5-10  $\mu$ g/ml induced significantly higher levels of T cell proliferation than did B7-H3Ig (Fig. 8A). In the absence of anti-CD3 mAb, neither B7-H3Ig nor B7-1Ig induced proliferation of T cells. Immobilized B7-H3Ig significantly enhanced proliferation of both CD4<sup>+</sup> and CD8<sup>+</sup> T cells (Fig. 8B).

To evaluate the ability of B7-H3 to affect CTL generation, purified human T cells from healthy donors were stimulated with cells of the melanoma line 624mel transiently transfected with pcDNA/B7-H3 or control vector. CTL activity was determined by lysis of <sup>51</sup>Cr-labeled 624mel cells. B7-H3/624mel cells induced significantly higher CTL activity than did mock-transfected 624mel cells (Fig. 8C). Thus B7-H3 co-stimulates the growth of both CD4+ and CD8+ T cells and enhances the generation of CTL.

#### Example 5. Molecular Cloning and Expression Pattern of B7-H4 encoding DNA

Using a strategy similar to that described above for B7-H3, the cDNA nucleotide sequence of an orf (SEQ ID NO:6) (Fig. 9) encoding another B7-homologue, B7-H4, was identified. This orf encodes a 282 amino acid protein (B7-H4; SEQ ID NO:5) (Fig. 10) which has significant homology in its predicted extracellular domain with human B7-H1, human B7-H2, human B7-H3, human B7-1, and human B7-2 (Fig. 11). The B7-H4 protein contains a typical signal peptide in its N-terminus, extracellular V- and C-like Ig domains, a transmembrane region, and a cytoplasmic tail, thereby indicating that B7-H4 is a type I transmembrane protein belonging to the Ig superfamily. Similar to the other members of the B7 family, B7-H4 has four conserved cysteine residues (marked \*) that are believed to be involved in the formation of V- and C-like Ig domains. The above data suggest that B7-H4 is a member of the B7 co-stimulatory ligand family.

Northern blot analysis was performed to determine mRNA expression of the B7-H4 gene. The analysis indicated that B7-H4 mRNA is expressed at relatively high levels in the lymphoid tissues spleen and thymus and weakly in lung.

#### Example 6. B7-H4 Co-stimulates T Cell Proliferation

In order to carry out T cell co-stimulation experiments, fusion proteins containing the extracellular domain of B7-H4 and the CH2-CH3 portion of either mouse IgG2a heavy chain (B7-H4Ig) or human IgG1 (B7-H4hIg) were prepared by recombinant methods substantially the same as that described above for the



production of B7-H3Ig. In the co-stimulation assays, nylon wool purified T cells were stimulated *in vitro* by immobilized anti-CD3 mAb in the presence of the immobilized fusion proteins containing the extracellular domains of the B7-H4 (B7-H4Ig or B7-H4hIg) or B7-1 (B7-1Ig). Control culture wells contained immobilized anti-CD3 mAb and control human IgG (hIgG) or contained immobilized anti-CD3 mAb and were "mock" coated with phosphate buffered saline (PBS) ("PBS"). The anti-CD3 mAb was coated onto the bottoms of microtiter plate tissue culture wells at the indicated concentrations (Fig. 12 and Fig. 13) and the fusion or control proteins were coated onto the bottoms of the wells at either 5 µg/ml (Fig. 12) or 10 µg/ml (Fig. 13). The assays were carried out as described above for the experiments testing for B7-H3 co-stimulatory activity with T cell proliferation being expressed in terms of the amount ("CPM") of [<sup>3</sup>H]-thymidine incorporated into the T cells after a 72 h culture. While coating at a concentration of 5 µg/ml resulted in a similar level of T cell co-stimulation with B7-H4Ig, B7-H4hIg, and B7-1Ig (Fig. 12), coating at a concentration of 10 µg/ml resulted in significantly higher T cell co-stimulation with B7-H4Ig and B7-H4hIg than with B7-1Ig (Fig. 13). These data indicate that B7-H4 co-stimulates the proliferation of T cells.

#### Example 7. B7-H3 Co-stimulates the Production Selectively of IFN-γ

Two different assays were used to test whether B7-H3 co-stimulates the production of a variety of cytokines. The first assay employed a cytokine mRNA array to measure the relative quantities of mRNA encoding 23 different cytokines in cells activated with anti-CD3 antibody in the presence of either control Ig, B7-1Ig, or B7-H3Ig (see above). The data are shown in Fig. 14. While the presence of B7-H3Ig resulted in detectable increases in mRNA encoding several cytokines (e.g., IL-1b, IL-2, IL-5, IL-6, IL-8, IL-12 (p35), IL-17, and TNF-α), it caused a remarkable increase in the level of IFN-γ mRNA.

A previously described co-stimulation assay was used [Dong et al. (1999) *Nat. Med.* 5, 1365-1369] to test whether B7-H3 co-stimulates the production of IFN-γ, IL-2 and IL-10 at the protein level. In this assay, purified T cells were stimulated *in vitro* by immobilized anti-CD3 mAb in the presence of immobilized control Ig, B7-

1Ig, or B7-H3Ig. After 72 h of culture, supernatants from all the cultures were removed and assayed for levels of the three cytokines by ELISA. Data obtained with T cells from four healthy human subjects are presented in Table 1; data from each individual and means  $\pm$  standard deviation are shown. While the presence of B7-H3Ig resulted in small but consistent increases in the level of IL-10, it resulted in dramatic increases in the level of IFN- $\gamma$ .

Thus it appears that B7-H3 selectively co-stimulates the production of IFN- $\gamma$  by T cells.

Table 1. Co-stimulation of cytokine production by B7-1Ig and B7-H3Ig

Co-stimulatory Polypeptide	Subject	Amount of Cytokine Produced		
		IL-2 (U/ml)	IL-10 (ng/ml)	IFN- $\gamma$ (ng/ml)
Control Ig	1	0.86	0.14	24.0
	2	0.87	0.13	37.75
	3	0.51	0.12	1.31
	4	0.39	0.07	7.72
	<u>Mean<math>\pm</math>SD</u>	0.66 $\pm$ 12	0.12 $\pm$ 0.02	17.70 $\pm$ 8.22
B7-1Ig	1	0.81	1.29	23.20
	2	0.72	1.63	142.40
	3	1.38	1.03	8.25
	4	1.67	0.44	8.50
	<u>Mean<math>\pm</math>SD</u>	1.15 $\pm$ 0.23	1.10 $\pm$ 0.25	0.87 $\pm$ 0.28
B7-H3Ig	1	0.53	0.64	26.61
	2	0.52	1.66	247.08
	3	0.89	0.84	12.58
	4	0.84	0.34	10.25
	<u>Mean<math>\pm</math>SD</u>	0.69 $\pm$ 0.10	0.87 $\pm$ 0.28	74.13 $\pm$ 57.76

Although the invention has been described with reference to the presently preferred embodiment, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims.

What is claimed is:

1. An isolated DNA comprising:
  - (a) a nucleic acid sequence that encodes a polypeptide with the ability to co-stimulate a T cell, wherein the nucleic acid sequence hybridizes under  
5 stringent conditions to the complement of a sequence that encodes a polypeptide with an amino acid sequence with SEQ ID NO:5; or
  - (b) the complement of the nucleic acid sequence.
2. The DNA of claim 1, wherein the nucleic acid sequence encodes a polypeptide comprising an amino acid sequence with SEQ ID NO:5.
- 10 3. The DNA of claim 1, wherein the nucleic acid sequence has a sequence of SEQ ID NO:6.
4. An isolated co-stimulatory polypeptide encoded by the DNA of claim 1.
5. The isolated polypeptide of claim 4, wherein the polypeptide comprises an amino acid sequence of amino acid residue 31 to amino acid residue 282  
15 of SEQ ID NO:5, or said amino acid sequence but with one or more conservative substitutions.
6. The isolated polypeptide of claim 5, wherein the polypeptide comprises an amino acid sequence of SEQ ID NO:5, or said amino acid sequence but with one or more conservative substitutions.
- 20 7. A vector comprising the DNA of claim 1.
8. The vector of claim 7, wherein the nucleic acid sequence is operably linked to a regulatory element which allows expression of said nucleic acid sequence in a cell.
9. A cell comprising the vector of claim 7.
- 25 10. A method of co-stimulating a T cell, the method comprising contacting the T cell with the polypeptide of claim 4.
11. The method of claim 10, wherein the contacting comprises culturing the polypeptide with the T cell *in vitro*.

12. The method of claim 10, wherein the T cell is in a mammal.
13. The method of claim 12, wherein the contacting comprises administering the polypeptide to the mammal.
14. The method of claim 12, wherein the contacting comprises  
5 administering a nucleic acid encoding the polypeptide to the mammal.
15. The method of claim 12, comprising:
- (a) providing a recombinant cell which is the progeny of a cell obtained from the mammal and has been transfected or transformed *ex vivo* with a nucleic acid encoding the polypeptide so that the cell expresses the polypeptide; and
  - 10 (b) administering the cell to the mammal.
16. The method of claim 15, wherein the recombinant cell is an antigen presenting cell (APC) and expresses the polypeptide on its surface.
17. The method of claim 16, wherein, prior to the administering, the APC is pulsed with an antigen or an antigenic peptide.
- 15 18. The method of claim 15, wherein the cell obtained from the mammal is a tumor cell.
19. The method of claim 12, wherein the mammal is suspected of having an immunodeficiency disease.
20. A method of identifying a compound that inhibits an immune response,  
20 the method comprising:
- (a) providing a test compound;
  - (b) culturing, together, the compound, the polypeptide of claim 4, a T cell, and a T cell activating stimulus; and
  - (c) determining whether the test compound inhibits the response of the T  
25 cell to the stimulus, as an indication that the test compound inhibits an immune response.
21. The method of claim 20, wherein the stimulus is an antibody that binds to a T cell receptor or a CD3 polypeptide.

22. The method of claim 20, wherein the stimulus is an alloantigen or an antigenic peptide bound to a major histocompatibility complex (MHC) molecule on the surface of an antigen presenting cell (APC).

23. The method of claim 22, wherein the APC is transfected or  
5 transformed with a nucleic acid encoding the polypeptide and the polypeptide is expressed on the surface of the APC.

24. A method of identifying a compound that enhances an immune response, the method comprising:

- (a) providing a test compound;
- 10 (b) culturing, together, the compound, the polypeptide of claim 4, a T cell, and a T cell activating stimulus; and
- (c) determining whether the test compound enhances the response of the T cell to the antigen, as an indication that the test compound enhances an immune response.

15 25. The method of claim 24, wherein the stimulus is an antibody that binds to a T cell receptor or a CD3 polypeptide.

26. The method of claim 25, wherein the stimulus is an alloantigen or an antigenic peptide bound to a MHC molecule on the surface of an APC.

27. The method of claim 26, wherein the APC is transfected or  
20 transformed with a nucleic acid encoding the polypeptide and the polypeptide is expressed on the surface of the APC.

28. An antibody that binds specifically to the polypeptide of claim 4.

29. The antibody of claim 28, wherein the antibody is a polyclonal antibody.

25 30. The antibody of claim 28, wherein the antibody is a monoclonal antibody.

31. The antibody of claim 28, wherein the antibody binds to the polypeptide with SEQ ID NO:5.

32. A cell comprising the vector of claim 8.

33. A method of producing a polypeptide that co-stimulates a T cell, the method comprising culturing the cell of claim 32 and purifying the polypeptide from the culture.

34. A fusion protein comprising a first domain joined to at least one additional domain, wherein the first domain comprises a polypeptide of claim 4.

35. The fusion protein of claim 34, wherein the at least one additional domain comprises the constant region of an immunoglobulin heavy chain or a fragment thereof.

36. A nucleic acid molecule encoding the fusion protein of claim 35.

37. A vector comprising the nucleic acid molecule of claim 36.

38. The vector of claim 37, wherein the nucleic acid molecule is operably linked to a regulatory element which allows expression of the nucleic acid molecule in a cell.

39. A cell comprising the vector of claim 38.

40. A method of producing a fusion protein, the method comprising culturing the cell of claim 39 and purifying the fusion protein from the culture.

41. A method of co-stimulating a T cell, the method comprising contacting the T cell with:

(a) a first co-stimulatory polypeptide selected from the group consisting of

(i) B7-H1, (ii) B7-H2, (iii) B7-H3, (iv) B7-H4, (v) a functional fragment of any of (i) - (iv), and (vi) any of (i) - (v) but with one or more conservative substitutions; and

(b) one or more additional co-stimulatory polypeptides selected from the group consisting of (vi) B7-1, (vii) B7-2, (viii) B7-H1, (ix) B7-H2, (x) B7-H3, (xi) B7-H4, (xii) a functional fragment of any of (vi) - (xi), and (xii) any of (vi) - (xii) but with one or more conservative substitutions.

42. The method of claim 41, wherein the contacting comprises culturing the first co-stimulatory polypeptide and the one or more additional co-stimulatory polypeptides with the T cell *in vitro*.

43. The method of claim 41, wherein the T cell is in a mammal.
44. The method of claim 43, wherein the contacting comprises administering the first co-stimulatory polypeptide and the one or more additional co-stimulatory polypeptides to the mammal.
- 5 45. The method of claim 43, wherein the contacting comprises administering one or more nucleic acids encoding the first co-stimulatory polypeptide and the one more additional co-stimulatory polypeptides to the mammal.
46. The method of claim 43, comprising:
- (a) providing a recombinant cell which is the progeny of a cell  
10 obtained from the mammal and which has been transfected or transformed *ex vivo* with one or more nucleic acids encoding the first co-stimulatory polypeptide and the one or more additional polypeptides so that the cell expresses the first co-stimulatory polypeptide and the one or more additional co-stimulatory polypeptides; and
- (b) administering the cell to the mammal.
- 15 47. The method of claim 43, comprising:
- (a) providing a first recombinant cell which is the progeny of a cell obtained from the mammal and which has been transfected or transformed *ex vivo* with a nucleic acid encoding the first co-stimulatory polypeptide;
- (b) providing one or more additional recombinant cells each of which  
20 is the progeny of a cell obtained from the mammal and each of which has been transfected or transformed *ex vivo* with a nucleic acid encoding one of the additional one or more co-stimulatory polypeptides; and
- (c) administering the first cell and the one or more additional cells to the mammal.
- 25 48. The method of claim 46, wherein the recombinant cell is an antigen presenting cell (APC) and expresses the first co-stimulatory polypeptide and the one or more additional co-stimulatory polypeptides on its surface.
49. The method of claim 48, wherein, prior to the administering, the APC is pulsed with an antigen or an antigenic peptide.



50. The method of claim 46, wherein the cell obtained from the mammal is a tumor cell.

51. The method of claim 43, wherein the mammal is suspected of having an immunodeficiency disease.

5 52. The method of claim 10, wherein the polypeptide co-stimulates the production of interferon- $\gamma$  by the T cell.

1/10

ATGCTGCGTCGGCGGGGAGCCCTGGCATGGGTGTGCATGTGGGTGCAGCCC  
TGGGAGCACTGTGGTTCTGCCTCACAGGAGCCCTGGAGGTCCAGGTCCCTGA  
AGACCCAGTGGTGGCACTGGTGGGCACCGATGCCACCCTGTGCTGCTCCTTCT  
CCCCTGAGCCTGGCTTCAGCCTGGCACAGCTCAACCTCATCTGGCAGCTGAC  
AGATACCAAACAGCTGGTGCACAGCTTTGCTGAGGGCCAGGACCAGGGCAG  
CGCCTATGCCAACCACGCGCCCTCTTCCCGACCTGCTGGCACAGGGCAAC  
GCATCCCTGAGGCTGCAGCGCGTGGTGTGGCGGACGAGGGCAGCTTACCT  
GCTTCGTGAGCATCCGGGATTTCCGCAGCGCTGCCGTACAGCTGCAGGTGGC  
CGCTCCCTACTCGAAGCCCAGCATGACCCTGGAGCCCAACAAGGACCTGCGG  
CCAGGGGACACGGTGACCATCACGTGCTCCAGCTACCGGGGCTACCCTGAGG  
CTGAGGTGTTCTGGCAGGATGGGCAGGGTGTGCCCCTGACTGGCAACGTGAC  
CACGTGCGAGATGGCCAACGAGCAGGGCTTGTGTTGATGTGCACAGCGTCTG  
CGGGTGGTGTGGGTGCGAATGGCACCTACAGCTGCCTGGTGCAGCAACCCCG  
TGCTGCAGCAGGATGCGCACGGCTCTGTCAACATCACAGGGCAGCCTATGAC  
ATTCCCCCAGAGGGCCCTGTGGGTGACCGTGGGGCTGTCTGTCTGTCTATTG  
CACTGCTGGTGGCCCTGGCTTTTCGTGTGCTGGAGAAAGATCAAACAGAGCTG  
TGAGGAGGAGAATGCAGGAGCTGAGGACCAGGATGGGGAGGGAGAAGGCTC  
CAAGACAGCCCTGCAGCCTCTGAAACACTCTGACAGCAAAGAAGATGATGG  
ACAAGAAATAGCCTGA

FIG. 1

ATGCTGCGTCGGCGGGGAGCCCTGGCATGGGTGTGCATGTGGGTGCAGCCC  
TGGGAGCACTGTGGTTCTGCCTCACAGGAGCCCTGGAGGTCCAGGTCCCTGA  
AGACCCAGTGGTGGCACTGGTGGGCACCGATGCCACCCTGTGCTGCTCCTTCT  
CCCCTGAGCCTGGCTTCAGCCTGGCACAGCTCAACCTCATCTGGCAGCTGAC  
AGATACCAAACAGCTGGTGCACAGCTTTGCTGAGGGCCAGGACCAGGGCAG  
CGCCTATGCCAACCACGCGCCCTCTTCCCGACCTGCTGGCACAGGGCAAC  
GCATCCCTGAGGCTGCAGCGCGTGGTGTGGCGGACGAGGGCAGCTTACCT  
GCTTCGTGAGCATCCGGGATTTCCGCAGCGCTGCCGTACAGCTGCAGGTGGC  
CGCTCCCTACTCGAAGCCCAGCATGACCCTGGAGCCCAACAAGGACCTGCGG  
CCAGGGGACACGGTGACCATCACGTGCCCCAGCTACCGGGGCTACCCTGAGG  
CTGAGGTGTTCTGGCAGGATGGGCAGGGTGTGCCCCTGACTGGCAACGTGAC  
CACGTGCGAGATGGCCAACGAGCAGGGCTTGTGTTGATGTGCACAGCGTCTG  
CGGGTGGTGTGGGTGCGAATGGCACCTACAGCTGCCTGGTGCAGCAACCCCG  
TGCTGCAGCAGGATGCGCACGGCTCTGTCAACATCACAGGGCAGCCTATGAC  
ATTCCCCCAGAGGGCCCTGTGGGTGACCGTGGGGCTGTCTGTCTGTCTATTG  
CACTGCTGGTGGCCCTGGCTTTTCGTGTGCTGGAGAAAGATCAAACAGAGCTG  
TGAGGAGGAGAATGCAGGAGCTGAGGACCAGGATGGGGAGGGAGAAGGCTC  
CAAGACAGCCCTGCAGCCTCTGAAACACTCTGACAGCAAAGAAGATGATGG  
ACAAGAAATAGCCTGA

FIG. 2

SUBSTITUTE SHEET (RULE 26)

2/10

MLRRRGSPGMGVHVGAALGALWFCLTGALEVQVPEDPVVALVGTDATLCCSFS  
 PEPGFSLAQLNLIWQLTDTKQLVHSFAEGDQGSAYANRTALFPDLLAQGNASL  
 RLQVRVVADEGSFTCFVSIRDFGSAAVSLQVAAPYSKPSMTLEPNKDLRPGDVT  
 ITCSSYRGYPEAEVFWQDGGQVPLTGNVTTSQMANEQGLFDVHSLRVVLGAN  
 GTYSCLVRNPVLQQDAHGSVTTTGQPMTFPPEALWVTVGLSVCLIALLLVALAFV  
 CWRKIKQSCEENAGAEDQDGEGEKSKTALQPLKHSDSKEDDGQEIA

FIG. 3

MLRRRGSPGMGVHVGAALGALWFCLTGALEVQVPEDPVVALVGTDATLCCSFS  
 PEPGFSLAQLNLIWQLTDTKQLVHSFAEGDQGSAYANRTALFPDLLAQGNASL  
 RLQVRVVADEGSFTCFVSIRDFGSAAVSLQVAAPYSKPSMTLEPNKDLRPGDVT  
 ITCPSYRGYPEAEVFWQDGGQVPLTGNVTTSQMANEQGLFDVHSLRVVLGAN  
 GTYSCLVRNPVLQQDAHGSVTTTGQPMTFPPEALWVTVGLSVCLIALLLVALAFV  
 CWRKIKQSCEENAGAEDQDGEGEKSKTALQPLKHSDSKEDDGQEIA

FIG. 4




FIG. 8A




FIG. 8B




FIG. 8C E:T

SUBSTITUTE SHEET (RULE 26)

h87-H3 1 MLRRRGSPGWHVGAA---LGAIFGLTGALEVOVPEE--HVAIVGTATLCOSESPEPGFSLAQINLIMQ 68  
h87-H2 1 MRLBS-----PGLFLIFS-SLRADTOEK--EMRAVVGSDVELSCACPEGRFDLNDIMYMQ 55  
h87-H1 1 MRTHAVFT---FMITYMHLIN-AFTVTVPKD--LYVVEYGSNMTTECKRPVEKQLDLAALITVWE 58  
h87-2 1 KGLSN-----ILRWAF-LLSGAPLK---ITGVFNEADLPDGFANSQNSLSLWVFMQ 52  
h87-1 1 MGHTRRQETSPSKCPYLNFEFLIMVAGLSHFCSGVTHMTKEKVAATLSQHNVS-VEELAQTRIMQ 67

h87-H3 69 LTD--TKDVRHSFAE--GDDGSAVARTALFPDLACGVASLRQRVRVADEGSHFDRS-IRDFESAAYS- 135  
h87-H2 56 TSEKTIWTVHIPQSSLENVDSRYNRALVSPAGLRGDSLRFLNMTBDEQKPHCLVL-SDSLGFQWLS 127  
h87-H1 59 MEDKNIIQFVHGEED--LKVOHSSYRORARLKQDLSLGMALQITDKLQDAGWRCMIS-YGGADYKRIT- 127  
h87-2 53 DQENLVNEMTGKEK-FDSVPSKMGKTSFD-----SDSMTLRHLNQLKOKQLVQCITHKKPTGVIRJHQ 119  
h87-1 68 KEK--KWLTHMSGD---MNLWPEYKRIITFD-----ITNLSTJMLAPPSDEGHECVVLKYEKDAFKREHL 131

h87-H3 136 LQVAPYSPKPSMTLEPNKOLPEE---TJTITCSSYRGYPEAE---VFDQGGGVPLTGNVITSQMANE---QG 201  
h87-H2 128 VEVLHMAVSVMAVMSAPHSPQD-ELTETCTSTNGYPEN---MMINKIDNSLDOALQNDTVFLNWRGL 196  
h87-H1 128 VRVAPYN---KINGRILVVDVTS-EHELTCGAL-EGYPAE---MILTSSDHQVLSGKTITNSKRE---EK 190  
h87-2 120 MNSLSMLANFSQPEMPSINJITENVYINLTCSSTHYPERKKMSMLRTKSTTIEYDGMQKSDNV---TEL 190  
h87-1 132 AEVTLNKADEPTPSJSDFELETJSN-IRRLTSTSGGFEPEH---LSALE-NGEEVAINITVSQDPE---TEL 197

h87-H3 202 FDVHSMRLMVL---GANEITSCLVRRPMLQO-----DAHGS-VITTEQPMITPPEALWVINGISVOLIA 261  
h87-H2 197 YDVSMALRIAR---TPSVNIGCCHEVILLOONLTVGSQTDGIGERDKLTENPVSTGEKN-AATKNSILAVLCL 265  
h87-H1 191 FVNTSLRINT---TTNELFYCTERRLDPEE-----NHTAE-LVPEELPLAPFNERTHLVLSAILQ 250  
h87-2 191 YDVSTSLSSFSFDPVTSMTLFCLETDKTL-----SSPFS--IELEDPPEPDHIPWITAVLPTVIG 253  
h87-1 198 YAVSSKLDFNM---ITVSEFCLIKYGLRY-----NOTFN---KNTTKQEHFBDNLLPSWATITJSMN 255

h87-H3 262 ILVALAFVC-PRRIKQSCFEENAGAEQDQSGEGSKTALQPLKPSDSKEDDGGQETIA 316  
h87-H2 266 LVWAVATG-WGDRQLHSYAGAWMSPEIT-----ELTGMV 302  
h87-H1 251 LGVALTFHIF-RLRGRWMDVKKCGEIDTNSKK-----QSDHLEET 290  
h87-2 264 VVWFCLILWKKKKRPNISYKGTINTMERES-EQTKKREKIHIPERSDEQRVFKSSKTSCKDSKDTCF 323  
h87-1 256 GUFVECDLIF-YCFAPRQERR--RNERLRRES-----VRPM 288

FIG. 5A

4/10

signal peptide  
 MLRRRGSPGMGVHVGAAALGALWFCLTGAL<sup>EVQVPEDPVVALVGTDA</sup>TLC<sup>CSFSPEPGFSLAQLNLINQLTDTKQLVHSFAEGDQGSAY</sup>  
 ANRTALFPDLLAQGNASLRQVRVAD<sup>EGSFTCFVSIRDFGSAAVSLQVAAPY</sup>SKPSMTLEPNKDLRPGDTVTIT<sup>CS</sup>SYRGYP<sup>EA</sup>EVFW  
 QDGGVPLTGNVTT<sup>SQMANEQGLFDVHSLRVVLGANGTYSCLVRNPVLQDDAHGSVTTTGQPMITFPPEALWTVGLSVCLIAL</sup>LVALA  
 FVCWRKIKQSC<sup>EEENAGAEDQDGE</sup>GSKTALQPLKHS<sup>DSKEDD</sup>QQEIA  
 IC

FIG. 5B

SUBSTITUTE SHEET (RULE 26)

5/10




FIG. 6A




FIG. 6B

SUBSTITUTE SHEET (RULE 26)

6/10




FIG. 7A




FIG. 7B

7/10

ATGGCTTCCCTGGGGCAGATCCTCTTCTGGAGCATAATTAGCATCATCATTAT  
TCTGGCTGGAGCAATTGCACTCATCATTGGCTTTGGTATTTTCAGGGAGACACT  
CCATCACAGTCACTACTGTGCGCTCAGCTGGGAACATTGGGGAGGATGGAAT  
CCTGAGCTGCACTTTGAACCTGACATCAAACCTTTCTGATATCGTGATACAAT  
GGCTGAAGGAAGGTGTTTTAGGCTTGGTCCATGAGTTCAAAGAAGGCAAAGA  
TGAGCTGTGCGAGCAGGATGAAATGTTTCAGAGGCCGGACAGCAGTGTTTGCT  
GATCAAGTGATAGTTGGCAATGCCTCTTTGCGGCTGAAAAACGTGCAACTCA  
CAGATGCTGGCACCTACAAATGTTATATCATCACTTCTAAAGGCAAGGGGAA  
TGCTAACCTTGAGTATAAACTGGAGCCTTCAGCATGCCGGAAGTGAATGTG  
GACTATAATGCCAGCTCAGAGACCTTGCGGTGTGAGGCTCCCCGATGGTTCC  
CCCAGCCACAGTGGTCTGGGCATCCCAAGTTGACCAGGGAGCCAACTTCTC  
GGAAGTCTCCAATACCAGCTTTGAGCTGAACTCTGAGAATGTGACCATGAAG  
GTTGTGTCTGTGCTCTACAATGTTACGATCAACAACACATACTCCTGTATGAT  
TGAAAATGACATTGCCAAAGCAACAGGGGATATCAAAGTGACAGAATCGGA  
GATCAAAAGGCGGAGTCACCTACAGCTGCTAAACTCAAAGGCTTCTCTGTGT  
GTCTCTTCTTTCTTTGCCATCAGCTGGGCACTTCTGCCTCTCAGCCCTTACCT  
GATGCTAAAATAA

FIG. 9

MASLGQILFWSIISIIILAGAIALIIGFGISGRHSITVTTVASAGNIGEDGILSCTFEPD  
IKLSDIVIQLKEGVLGLVHEFKEGKDELSEQDEMFRGRTAVFADQVIVGNASLR  
LKNVQLTDAGTYKCYIITSKGKGNANLEYKTGAFSMPENVVDYNASSETLRCEA  
PRWFPQPTVVWASQVDQGANFSEVSNTSFELNSENVTMKVVSVLNVNTINNTYS  
CMIENDIAKATGDIKVTESIKRRSHLQLLNSKASLCVSSFFAISWALLPLSPYLM  
LK

FIG. 10



8/10

B7-H4 ISGRHSITVTIVASAGNITGEGITISOTIF--EPDIKISUTIMICOMKEG--VIGLVIHFRKEGKD  
 B7-H2 LRAOTQEKEVRAMGSDVIELSCACPEGRSFDINWYMYQTSSEKTMATYHTPONSSSL  
 B7-H3.2 QVPEDPVVALMGTDAITLCCSPSPGFSIQLNINQJLTD--TKQLMHSFAEGQD  
 B7-H1 VTVPKOLYVVEYSGNMTTECKTPVEKQLDIWALIVYMEMEDKNITQFMHGEEDIKV  
 B7-2 QAYFNETADLPQCFANSONQSISELWFMQDOENLVNEMYLGEKFD  
 B7-1 LNFFQLLVLAGLSHFCSGVIHVTKEKEVATISGHN-VSVEELATRIYHCKEK--KAVLTMMSCMNI

B7-H4 ELSEQDEMFGRITAMFADQIMVGNASRLKNVQITDAGITKQJITTSKKG-GNANLEYKTGAFS-----M  
 B7-H2 EN--VDSRYRNRAIMSPAGMLREDFSLRLFNITPODEQFHGLVLSQSLG-FOEMLSIEVILHVAANFSV  
 B7-H3.2 ----QGSANVARTALFFDLACGNASRLDRVRADERSHDEVSIRDFG-SAAVSLQVADPYSKPSMTL  
 B7-H1 ----QHSSVECRARLLKQLSLENAALQITDYKLDQAGVRCMISYGGAD-YKRIITVKVNPYN----KIN  
 B7-2 S---VHSKMGXTSFDSD----SWITRLNHLQIKOKGLVQCIIHHKPTGMIRIHOHNSSELSVLANFSQ  
 B7-1 -----WPEIKNRTIIF--D--ITMISIMIALRPSDEGTYECVYLKYEKDAFKREHLAEISVKADEFPT

B7-H4 PEIN--VDYNAS-SETLRCEAPRMFPQR-TMVASQVDIGANFSEVSNISFELSENITMKVWSMLY---  
 B7-H2 PMSAPHSPSOD-ELHETCISITNGYPRP-NMINKTNSLLDOALQNTIVELMIRGLYDWSMIRI  
 B7-H3.2 EPNKD-LRPGD--TVITTCPSYRGYPEA-EMFAD--GGVPLTGNVITTSQVANEQGLFDMHSWARM  
 B7-H1 ORILV-VDBVTS-EHELTCDA-EGYPKA-EMVATS--SHQVLSGKTITINSKREEKLFNWTSTLRIL  
 B7-2 PETVPISNITENWYINLTCSIIHGYPEPKVSVLLRTQNSTIYDGINOKSQNVTELVDNISIISUSFP  
 B7-1 PSUSDFEITISN-IRRTDSTISGEPPEP-HLSALE--NGEELNAINITVSDPETELYAUSSKLDLDF----

B7-H4 NVTIINTYSWITJENDIAKATGDI-----KVTESEIKRBSHLQLLSK  
 B7-H2 ARTIPSWNIGCJENWILLQNLVNGSQTGNDIGERDKJENFVSTIGEKNAT  
 B7-H3.2 VILGANITYSCLVRNPVLDDAHG-----SVTITGCP-MTFPI  
 B7-H1 NITITNEIYQIFRRLDPEENITA-----ELVIPELP-LAHPPNERTH  
 B7-2 DVTSNMIFCTILEIDKTRLLSSP-----FSIELEDD--QPPPDHIP  
 B7-1 NMTIN-SEMLJIKYGHLRVNDIF-----NWNITIKOE--HFPDN

FIG. 11

9/10




FIG. 12




FIG. 13

SUBSTITUTE SHEET (RULE 26)

10/10




FIG. 14

SUBSTITUTE SHEET (RULE 26)

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/41430

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07H 21/04; C12N 15/00, 15/12, 15/63; C07K 16/46

US CL : 526/23.4, 23.5; 435/69.1, 320.1, 252.3; 530/387.3

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 526/23.4, 23.5; 435/69.1, 320.1, 252.3; 530/387.3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Please See Continuation Sheet

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO99/63088 A1 (GENENTECH INC.) 09 December 1999 (09.12.1999), pages 1-2, 27-28, 188-190, 280-284, 294, 334-335, 451, 497-500 and Figures 207 and 208 with respect to PRO1291.	1-3, 7-9, 32-40

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"B" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z"

document member of the same patent family

Date of the actual completion of the international search

18 November 2001 (18.11.2001)

Date of mailing of the international search report

27 DEC 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Jessica H. Roark

Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/41430

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claim Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:  
Please See Continuation Sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1-3, 7-9 and 32-40

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/41430

### BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

1. This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claims 1-3, 7-9, 32-40, drawn to DNAs related to SEQ ID NO:6 encoding SEQ ID NO:5 as well as fusion protein of SEQ ID NO:5 and the DNA encoding; vectors, host cells and methods of producing the polypeptide and fusion protein.

Group II, claims 4-6, drawn to co-stimulatory polypeptides related to SEQ ID NO:5.

Group III, claims 10-13, 19 and 52, drawn to a method of co-stimulating a T cell by administering a polypeptide related to SEQ ID NO:5.

Group IV, claims 10, 12 and 14-19, drawn to a method of co-stimulating a T cell by administering a nucleic acid encoding a polypeptide related to SEQ ID NO:5.

Group V, claims 20-23, drawn to a method of identifying a compound that inhibit an immune response.

Group VI, claims 24-27, drawn to a method of identifying a compound that enhances an immune response.

Group VII, claims 28-31, drawn to an antibody to a polypeptide related to SEQ ID NO:5.

Group VIII, claims 41-44 and 51, drawn to a method of co-stimulating a T cell comprising administering a first co-stimulatory polypeptide and one or more additional co-stimulatory polypeptides.

Group IX, claims 41, 43 and 45-51, drawn to a method of co-stimulating a T cell comprising administering one or more nucleic acids encoding a first co-stimulatory polypeptide and one or more additional co-stimulatory polypeptides.

2. This application contains claims directed to more than one species of the generic invention. These species are deemed to lack unity of invention because they are not so linked as to form a single general inventive concept under PCT Rule 13.1.

In order for more than one species to be examined, the appropriate additional examination fees must be paid. The species are as follows:

in Groups V and VI, wherein the stimulus is either an antibody or an APC;

in Group VIII, an ultimate species identifying the particular combination of the several co-stimulatory molecules.

The following claims are generic: in Group V, claim 20 is generic; in Group VI claim 24 is generic; and in Group VIII, claim 41 is generic.

3. The inventions listed as Groups I-IX and species therein do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The special technical feature is considered to be the nucleic acid of SEQ ID NO:6 which encodes SEQ ID NO:5. However, the nucleic acid of SEQ ID NO:6 fails to define a contribution over the prior art of WO99/63088. WO99/63088 teaches the DNA encoding the membrane bound protein PRO1291, which is identical to the full length sequence of instant SEQ ID NO:6.

**INTERNATIONAL SEARCH REPORT**

International application No.

PCT/US01/41430

**Continuation of B. FIELDS SEARCHED Item 3:**

Databases: Geneseq, WEST, Medline, Embase, Biosis

Search terms: SEQ ID NO:5, SEQ ID NO:6, inventor name, B7?

**THIS PAGE BLANK (USPTO)**